



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:39 PM GMT

PDB ID : 4YUV  
Title : Crystal structure of Trypanosoma cruzi spermidine synthase in complex with decarboxylated S-adenosylmethionine  
Authors : Amano, Y.; Tateishi, Y.  
Deposited on : 2015-03-19  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	
1	B	304	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Spermidine synthase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	0	0
			2233	1418	382	419	14			
1	B	294	Total	C	N	O	S	0	0	0
			2306	1464	393	434	15			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP Q4DA73
A	-6	ALA	-	expression tag	UNP Q4DA73
A	-5	HIS	-	expression tag	UNP Q4DA73
A	-4	HIS	-	expression tag	UNP Q4DA73
A	-3	HIS	-	expression tag	UNP Q4DA73
A	-2	HIS	-	expression tag	UNP Q4DA73
A	-1	HIS	-	expression tag	UNP Q4DA73
A	0	HIS	-	expression tag	UNP Q4DA73
B	-7	MET	-	initiating methionine	UNP Q4DA73
B	-6	ALA	-	expression tag	UNP Q4DA73
B	-5	HIS	-	expression tag	UNP Q4DA73
B	-4	HIS	-	expression tag	UNP Q4DA73
B	-3	HIS	-	expression tag	UNP Q4DA73
B	-2	HIS	-	expression tag	UNP Q4DA73
B	-1	HIS	-	expression tag	UNP Q4DA73
B	0	HIS	-	expression tag	UNP Q4DA73

- Molecule 2 is 5'-[(S)-(3-AMINOPROPYL)(METHYL)-LAMBDA 4 -SULFANYL]-5'-DEOXYADENOSINE (three-letter code: S4M) (formula: C<sub>14</sub>H<sub>24</sub>N<sub>6</sub>O<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			24	14	6	3	1		
2	B	1	Total	C	N	O	S	0	0
			24	14	6	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	133	Total	O	0	0
			133	133		
3	B	144	Total	O	0	0
			144	144		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Spermidine synthase, putative



- Molecule 1: Spermidine synthase, putative



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.35Å 92.36Å 68.27Å 90.00° 99.76° 90.00°	Depositor
Resolution (Å)	24.71 – 1.60 24.71 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.8 (24.71-1.60) 96.8 (24.71-1.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.84 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.196 , 0.240 0.195 , 0.240	Depositor DCC
$R_{free}$ test set	3508 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.9	Xtriage
Anisotropy	0.084	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.47 , 54.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 69054 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4864	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: S4M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.15	5/2287 (0.2%)	1.20	17/3099 (0.5%)
1	B	1.13	6/2364 (0.3%)	1.20	17/3207 (0.5%)
All	All	1.14	11/4651 (0.2%)	1.20	34/6306 (0.5%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	137	SER	CB-OG	8.46	1.53	1.42
1	B	162	TYR	CE2-CZ	6.09	1.46	1.38
1	B	53	TRP	CE3-CZ3	5.90	1.48	1.38
1	A	137	SER	CA-CB	5.72	1.61	1.52
1	B	274	TYR	CE1-CZ	5.65	1.45	1.38
1	B	274	TYR	CG-CD2	5.63	1.46	1.39
1	A	14	GLU	CG-CD	5.55	1.60	1.51
1	B	273	TYR	CG-CD2	5.52	1.46	1.39
1	A	274	TYR	CG-CD2	5.42	1.46	1.39
1	A	198	CYS	CA-CB	5.32	1.65	1.53
1	B	81	SER	CB-OG	5.07	1.48	1.42

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	103	ARG	NE-CZ-NH2	-9.39	115.60	120.30
1	A	190	ARG	NE-CZ-NH1	-9.10	115.75	120.30
1	B	260	ARG	NE-CZ-NH2	-8.62	115.99	120.30
1	B	65	VAL	CG1-CB-CG2	8.57	124.61	110.90
1	A	137	SER	N-CA-CB	-8.06	98.41	110.50
1	B	159	ASP	CB-CG-OD1	7.76	125.28	118.30
1	A	195	ASP	CB-CG-OD2	-6.92	112.07	118.30
1	A	140	ASP	CB-CG-OD2	-6.72	112.25	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	70	GLU	OE1-CD-OE2	-6.71	115.25	123.30
1	B	103	ARG	NE-CZ-NH1	6.50	123.55	120.30
1	B	120	ASP	CB-CG-OD2	-6.42	112.52	118.30
1	B	149	ASP	CB-CG-OD1	6.37	124.03	118.30
1	A	118	ASP	CB-CG-OD1	6.29	123.96	118.30
1	B	208	ASP	CB-CG-OD2	6.26	123.93	118.30
1	A	195	ASP	CB-CG-OD1	6.17	123.86	118.30
1	A	142	ARG	NE-CZ-NH1	-6.06	117.27	120.30
1	A	208	ASP	CB-CG-OD1	-5.96	112.93	118.30
1	B	260	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	B	209	LEU	CB-CG-CD2	-5.83	101.09	111.00
1	A	65	VAL	CG1-CB-CG2	5.82	120.20	110.90
1	B	142	ARG	NE-CZ-NH1	-5.81	117.40	120.30
1	A	65	VAL	CA-CB-CG1	5.80	119.61	110.90
1	A	82	LEU	CB-CG-CD2	-5.58	101.52	111.00
1	A	14	GLU	OE1-CD-OE2	-5.53	116.66	123.30
1	A	190	ARG	NE-CZ-NH2	5.51	123.06	120.30
1	A	65	VAL	CA-CB-CG2	5.45	119.08	110.90
1	B	155	ARG	CB-CG-CD	-5.40	97.57	111.60
1	B	140	ASP	CB-CG-OD1	5.33	123.10	118.30
1	B	224	PHE	CB-CG-CD1	5.29	124.50	120.80
1	B	70	GLU	OE1-CD-OE2	-5.23	117.03	123.30
1	B	65	VAL	CA-CB-CG2	5.19	118.69	110.90
1	B	98	ASP	CB-CG-OD1	-5.05	113.75	118.30
1	A	138	LEU	CB-CG-CD1	-5.05	102.42	111.00
1	A	142	ARG	NE-CZ-NH2	5.01	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2233	0	2191	30	0
1	B	2306	0	2257	17	0
2	A	24	0	24	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	24	0	24	0	0
3	A	133	0	0	6	0
3	B	144	0	0	4	0
All	All	4864	0	4496	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:ARG:HG3	3:B:729:HOH:O	1.58	1.02
1:A:290:ARG:HH11	1:A:290:ARG:HG2	1.22	1.01
1:A:90:ARG:NH2	3:A:601:HOH:O	1.92	1.00
1:B:1:MET:HB3	1:B:2:PRO:HD3	1.50	0.93
1:A:155:ARG:HG2	3:A:628:HOH:O	1.73	0.88
1:A:210:GLU:H	1:A:210:GLU:CD	1.98	0.67
1:A:290:ARG:NH1	1:A:290:ARG:HG2	2.00	0.65
1:B:182:GLU:HG3	1:B:218:PHE:CG	2.32	0.64
1:A:155:ARG:CG	3:A:628:HOH:O	2.38	0.64
1:A:152:ALA:O	1:A:156:GLN:HG3	1.97	0.64
1:B:5:GLU:CD	3:B:602:HOH:O	2.35	0.63
1:B:155:ARG:CG	3:B:729:HOH:O	2.31	0.61
1:B:214:LYS:HE2	1:B:217:ARG:HH22	1.67	0.60
1:A:13:ARG:NH1	1:B:11:TRP:CD2	2.71	0.58
1:A:179:LEU:N	1:A:179:LEU:HD12	2.19	0.57
1:B:1:MET:CB	1:B:2:PRO:HD3	2.29	0.56
1:A:290:ARG:HH11	1:A:290:ARG:CG	2.08	0.55
1:A:256:THR:O	1:A:280:LYS:HD3	2.07	0.53
1:A:64:GLN:OE1	2:A:501:S4M:HG1	2.10	0.51
1:A:176:ALA:HA	1:A:179:LEU:HD13	1.92	0.51
1:A:229:TYR:CZ	1:A:243:GLY:HA3	2.46	0.51
1:B:19:TRP:HB3	1:B:207:LEU:HD13	1.93	0.50
1:B:229:TYR:CZ	1:B:243:GLY:HA3	2.47	0.50
1:A:265:MET:HE2	3:A:712:HOH:O	2.14	0.47
1:A:219:ILE:O	1:A:222:THR:HG22	2.15	0.46
1:A:22:GLN:HE21	1:B:53:TRP:HE1	1.62	0.46
1:A:171:ASP:HB2	1:A:172:PRO:CD	2.46	0.45
1:A:57:MET:HB3	1:A:65:VAL:HG12	1.98	0.45
1:A:13:ARG:NH1	1:B:11:TRP:CE3	2.84	0.45
1:B:57:MET:HB3	1:B:65:VAL:HG12	1.99	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:HIS:HE1	3:B:731:HOH:O	2.00	0.44
1:B:182:GLU:HG3	1:B:218:PHE:CD1	2.52	0.44
1:A:210:GLU:O	1:A:213:GLU:HG2	2.18	0.44
1:B:213:GLU:HG3	1:B:292:ILE:HD11	1.99	0.44
1:A:128:LYS:HE2	3:A:713:HOH:O	2.17	0.43
1:A:210:GLU:CD	1:A:210:GLU:N	2.71	0.43
1:A:262:VAL:O	1:A:268:ALA:HB2	2.19	0.42
1:A:136:ARG:O	3:A:602:HOH:O	2.22	0.42
1:A:44:ILE:CD1	1:A:126:GLN:HB3	2.50	0.42
1:A:7:ILE:HG13	1:A:8:SER:N	2.35	0.42
1:B:189:LEU:HD22	1:B:222:THR:HG23	2.02	0.41
1:B:293:ASN:O	1:B:294:ASN:HB2	2.21	0.41
1:A:92:LEU:HD21	1:A:94:ILE:HD11	2.02	0.41
1:A:179:LEU:CD1	1:A:179:LEU:N	2.83	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/304 (92%)	271 (96%)	10 (4%)	0	100	100
1	B	292/304 (96%)	280 (96%)	12 (4%)	0	100	100
All	All	573/608 (94%)	551 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/262 (94%)	235 (96%)	10 (4%)	37	11
1	B	253/262 (97%)	246 (97%)	7 (3%)	51	22
All	All	498/524 (95%)	481 (97%)	17 (3%)	44	16

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	VAL
1	A	124	MET
1	A	182	GLU
1	A	186	LYS
1	A	195	ASP
1	A	210	GLU
1	A	218	PHE
1	A	222	THR
1	A	261	PRO
1	A	290	ARG
1	B	30	LYS
1	B	50	LYS
1	B	98	ASP
1	B	155	ARG
1	B	201	GLN
1	B	218	PHE
1	B	222	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	22	GLN
1	A	201	GLN
1	A	233	HIS
1	B	22	GLN
1	B	233	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	S4M	A	501	-	23,26,26	1.37	3 (13%)	17,37,37	2.07	4 (23%)
2	S4M	B	501	-	23,26,26	1.29	4 (17%)	17,37,37	2.25	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	S4M	A	501	-	-	0/7/28/28	0/3/3/3
2	S4M	B	501	-	-	0/7/28/28	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	S4M	C8-N7	-2.17	1.30	1.34
2	A	501	S4M	C4-N3	-2.02	1.32	1.35
2	B	501	S4M	C2-N3	2.02	1.35	1.32
2	B	501	S4M	O2'-C2'	2.42	1.48	1.43
2	B	501	S4M	O4'-C1'	2.52	1.44	1.41
2	A	501	S4M	C2'-C1'	2.56	1.57	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	S4M	CG-SD	3.46	1.88	1.80

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	S4M	C4'-O4'-C1'	-7.85	101.32	109.64
2	A	501	S4M	N3-C2-N1	-5.33	124.69	128.87
2	A	501	S4M	C4'-O4'-C1'	-3.76	105.66	109.64
2	B	501	S4M	N3-C2-N1	-3.17	126.38	128.87
2	A	501	S4M	C1'-N9-C4	-2.77	123.72	126.81
2	A	501	S4M	O4'-C1'-N9	3.37	114.48	108.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	S4M	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	285/304 (93%)	0.05	9 (3%) 51 48	11, 20, 34, 55	0
1	B	294/304 (96%)	0.10	13 (4%) 38 34	11, 19, 39, 58	0
All	All	579/608 (95%)	0.08	22 (3%) 44 41	11, 19, 37, 58	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	5.7
1	B	9	GLY	5.5
1	A	16	ASN	4.8
1	A	173	ALA	3.4
1	B	294	ASN	3.2
1	B	3	GLY	3.1
1	B	173	ALA	3.0
1	B	19	TRP	3.0
1	A	50	LYS	3.0
1	B	136	ARG	2.7
1	B	8	SER	2.6
1	A	20	PRO	2.5
1	B	5	GLU	2.4
1	A	166	ILE	2.3
1	A	136	ARG	2.3
1	A	251	ALA	2.3
1	A	49	PRO	2.3
1	B	183	ALA	2.3
1	B	49	PRO	2.2
1	B	7	ILE	2.1
1	B	6	LEU	2.1
1	A	294	ASN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	S4M	B	501	24/24	0.95	0.10	0.07	14,17,23,26	0
2	S4M	A	501	24/24	0.96	0.08	-0.55	12,14,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.