wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 06:58 PM BST

PDB ID	:	1YX8
Title	:	NMR structure of Calsensin, 20 low energy structures.
Authors	:	Venkitaramani, D.V.; Fulton, D.B.; Andreotti, A.H.; Johansen, K.M.; Johansen, J.
Deposited on	:	2005-02-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 81%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	83	6% 53% 31% 10%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 18 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:6-A:80 (75)	0.62	18

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 8 single-model clusters were found.

Cluster number	Models
1	4, 7, 11, 14, 16, 20
2	1, 6, 10, 18
3	3, 12
Single-model clusters	2; 5; 8; 9; 13; 15; 17; 19

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1301 atoms, of which 666 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Calsensin.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	83	1301	402	666	103	125	5	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Calsensin

Chain A: 6% 53% 31% 10%

M1 A2 C3 K4 V5 K6 L9 E10 A11 A12 F13 K14 K15 L16 D17 A18 N19 G20 D21 G22 Y23 V24 T25 A26 L27 E28 L29 Q30 T31 F32 M33 V34 T35 L36 D37 A38 Y39 K40 A41 L42 S43 K44 D45 K46 V47 K48 E49 A50 S51 A52 K53 L54 I55 K56 M57 A58 D59 K60 N61

S62 D63 G64 K65 I66 S67 K68 E69 E70 F71 L72 N73 A74 N75 A76 E77 L78 L79 C80 Q81 L82 K83

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 18. Colouring as in section 4.1 above.

- Molecule 1: Calsensin

Chain A: 7% 46% 37% 10%

M1 A2 C3 K4 V5 K6 L9 E10 A11 A12 F13 K14 K15 L16 D17 A18 N19 G20 D21 G22 Y23 V24 T25 A26 L27 E28 L29 Q30 T31 F32 M33 V34 T35 L36 D37 A38 Y39 K40 A41 L42 S43 K44 D45 K46 V47 K48 S51 A52 K53 L54 I55 K56 M57 A58 D59 K60 N61 S62

D63 G64 K65 I66 S67 K68 E69 E70 F71 L72 N73 A74 N75 A76 E77 L78 L79 C80 Q81 L82 K83

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *Distance geometry and simulated annealing.*

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with favorable non-bond energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6519
Number of chemical shift lists	1
Total number of shifts	763
Number of shifts mapped to atoms	763
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	81%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	573	591	591	122±5
All	All	11460	11820	11820	2439

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 105.

5 of 586 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:16:LEU:HD12	1:A:17:ASP:N	1.14	1.55	7	19
1:A:29:LEU:HD11	1:A:72:LEU:HD11	1.10	1.17	19	15
1:A:24:VAL:HG11	1:A:58:ALA:HB2	1.03	1.20	12	3
1:A:16:LEU:O	1:A:25:THR:HB	1.01	1.55	9	5
1:A:16:LEU:HD13	1:A:26:ALA:HB3	0.97	1.30	9	2

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	75/83 (90%)	54±2 (72±2%)	13±2 (18±3%)	8±2 (11±3%)	1 9
All	All	1500/1660 (90%)	1073 (72%)	266 (18%)	161 (11%)	1 9

5 of 20 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	20	GLY	20
1	A	67	SER	20
1	A	62	SER	18
1	A	80	CYS	13
1	A	22	GLY	13

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	60/67 (90%)	31±2 (51±4%)	29±2 (49±4%)	0 1
All	All	1200/1340 (90%)	617 (51%)	583 (49%)	0 1

5 of 54 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	48	LYS	20
1	A	9	LEU	20
1	A	72	LEU	20
1	A	53	LYS	20
1	A	40	LYS	20

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 81% for the well-defined parts and 73% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 6519

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	763
Number of shifts mapped to atoms	763
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	12

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	75	-0.86 ± 0.16	Should be applied
$^{13}C_{\beta}$	72	-0.59 ± 0.15	Should be applied
$^{13}C'$	0	—	—
^{15}N	74	-1.19 ± 0.26	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 81%, i.e. 723 atoms were assigned a chemical shift out of a possible 893. 4 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	297/375 (79%)	148/150 (99%)	75/150 (50%)	74/75 (99%)
Sidechain	403/475 (85%)	259/275 (94%)	139/183 (76%)	5/17 (29%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	23/43 (53%)	23/23 (100%)	0/20 (0%)	0/0 (—%)
Overall	723/893 (81%)	430/448 (96%)	214/353 (61%)	79/92 (86%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	57	MET	CE	42.56	26.97 – 7.37	13.0
1	A	54	LEU	CG	37.52	32.55 – 21.05	9.3
1	A	60	LYS	CG	35.12	30.67 – 19.17	8.9
1	A	78	LEU	CB	30.19	51.69 – 32.89	-6.4
1	A	15	LYS	HE2	4.08	3.87 – 1.97	6.1
1	A	75	ASN	HB3	4.77	4.41 – 1.11	6.1
1	A	44	LYS	HE2	4.06	3.87 – 1.97	6.0
1	A	49	GLU	HG2	1.06	3.33 – 1.23	-5.8
1	A	44	LYS	HE3	4.01	3.86 – 1.96	5.8
1	A	57	MET	CG	25.29	38.33 – 25.73	-5.3
1	A	60	LYS	HE3	1.90	3.86 – 1.96	-5.3
1	A	65	LYS	HE2	1.94	3.87 – 1.97	-5.1

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: