



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 21, 2016 – 07:27 PM EDT

PDB ID : 4Z0W  
Title : Peptaibol gichigamin isolated from Tolypocladium sup\_5  
Authors : Du, L.; Risinger, A.L.; Mitchell, C.A.; Stamps, B.W.; Pan, N.; King, J.B.;  
Motley, J.L.; Thomas, L.M.; Yang, Z.; Stevenson, B.S.; Mooberry, S.L.; Ci-  
chewicz, R.H.  
Deposited on : 2015-03-26  
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0122  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1006 (1.14-1.06)
Clashscore	102246	1055 (1.14-1.06)
Ramachandran outliers	100387	1016 (1.14-1.06)
Sidechain outliers	100360	1014 (1.14-1.06)
RSRZ outliers	91569	1009 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	23	<div> <div>39%</div> <div> <div></div> <div>70%</div> <div>22%</div> <div>.</div> <div>.</div> </div> </div>
1	B	23	<div> <div>43%</div> <div> <div></div> <div>61%</div> <div>30%</div> <div>9%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 608 atoms, of which 317 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEPTAIBOL GICHIGAMIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	22	Total	C	H	N	O	0	0	0
			294	96	156	21	21			
1	B	23	Total	C	H	N	O	0	1	0
			305	98	161	22	24			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	O	0	0
			2	2		
2	B	7	Total	O	0	0
			7	7		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### ● Molecule 1: PEPTAIBOL GICHIGAMIN



#### ● Molecule 1: PEPTAIBOL GICHIGAMIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	10.06Å 18.79Å 31.42Å 75.16° 88.54° 86.48°	Depositor
Resolution (Å)	25.00 – 1.10 18.13 – 1.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (25.00-1.10) 89.1 (18.13-1.10)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.10Å)	Xtriage
Refinement program	SHELXL	Depositor
R, $R_{free}$	0.110 , 0.121 0.134 , 0.138	Depositor DCC
$R_{free}$ test set	425 reflections (5.10%)	DCC
Wilson B-factor (Å <sup>2</sup> )	2.9	Xtriage
Anisotropy	0.703	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
Estimated twinning fraction	0.000 for -h,-k,-k+l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 9015 reflections	Xtriage
$F_o, F_c$ correlation	0.34	EDS
Total number of atoms	608	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BAL, DIV, ACE, AIB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.16	1/66 (1.5%)	1.75	1/86 (1.2%)
1	B	2.00	2/72 (2.8%)	2.04	4/88 (4.5%)
All	All	1.65	3/138 (2.2%)	1.91	5/174 (2.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	4
All	All	0	9

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	23[A]	GLY	C-OXT	9.55	1.41	1.23
1	B	23[B]	GLY	C-OXT	9.55	1.41	1.23
1	A	8	PRO	CA-C	-5.10	1.42	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	21	LEU	CA-CB-CG	5.65	128.29	115.30
1	A	3	PRO	N-CA-CB	5.45	109.83	103.30
1	B	3	PRO	N-CA-CB	5.42	109.80	103.30
1	B	6	PHE	CB-CG-CD2	5.36	124.55	120.80
1	B	8	PRO	N-CA-CB	5.08	109.40	103.30

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	AIB	Peptide
1	A	13	DIV	Peptide
1	A	16	AIB	Peptide
1	A	21	LEU	Peptide
1	A	3	PRO	Mainchain
1	B	10	AIB	Peptide
1	B	13	DIV	Peptide
1	B	16	AIB	Peptide
1	B	3	PRO	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	138	156	150	0	0
1	B	144	161	150	1	0
2	A	2	0	0	0	0
2	B	7	0	0	0	0
All	All	291	317	300	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:21:LEU:O	1:B:22:BA1:HA2	2.15	0.46

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	9/23 (39%)	8 (89%)	1 (11%)	0	100	100
1	B	9/23 (39%)	9 (100%)	0	0	100	100
All	All	18/46 (39%)	17 (94%)	1 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	7/7 (100%)	7 (100%)	0	100	100
1	B	7/7 (100%)	7 (100%)	0	100	100
All	All	14/14 (100%)	14 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

24 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	AIB	A	10	1	2,5,6	0.91	0	4,7,9	2.22	2 (50%)
1	BAL	A	11	1	4,4,5	0.84	0	1,3,5	1.13	0
1	DIV	A	13	1	3,6,7	0.82	0	3,8,10	1.42	1 (33%)
1	BAL	A	14	1	4,4,5	0.70	0	1,3,5	1.13	0
1	AIB	A	16	1	2,5,6	0.94	0	4,7,9	1.22	1 (25%)
1	BAL	A	17	1	4,4,5	0.88	0	1,3,5	0.84	0
1	AIB	A	19	1	2,5,6	0.98	0	4,7,9	1.10	1 (25%)
1	AIB	A	2	1	2,5,6	0.58	0	4,7,9	2.02	1 (25%)
1	AIB	A	20	1	2,5,6	0.73	0	4,7,9	1.13	1 (25%)
1	BAL	A	22	1	3,3,5	0.17	0	0,2,5	0.00	-
1	AIB	A	4	1	2,5,6	1.22	0	4,7,9	1.66	1 (25%)
1	DIV	A	7	1	3,6,7	1.13	0	3,8,10	1.77	1 (33%)
1	AIB	B	10	1	2,5,6	1.07	0	4,7,9	1.78	1 (25%)
1	BAL	B	11	1	4,4,5	0.44	0	1,3,5	1.14	0
1	DIV	B	13	1	3,6,7	0.54	0	3,8,10	1.30	1 (33%)
1	BAL	B	14	1	4,4,5	0.72	0	1,3,5	1.66	0
1	AIB	B	16	1	2,5,6	0.91	0	4,7,9	0.80	0
1	BAL	B	17	1	4,4,5	0.59	0	1,3,5	1.10	0
1	AIB	B	19	1	2,5,6	0.89	0	4,7,9	1.42	1 (25%)
1	AIB	B	2	1	2,5,6	0.74	0	4,7,9	1.78	2 (50%)
1	AIB	B	20	1	2,5,6	0.83	0	4,7,9	1.55	1 (25%)
1	BAL	B	22	1	4,4,5	1.03	0	1,3,5	0.03	0
1	AIB	B	4	1	2,5,6	1.15	0	4,7,9	2.53	2 (50%)
1	DIV	B	7	1	3,6,7	1.25	0	3,8,10	1.32	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	AIB	A	10	1	-	0/2/3/6	0/0/0/0
1	BAL	A	11	1	-	0/1/2/3	0/0/0/0
1	DIV	A	13	1	-	0/3/6/9	0/0/0/0
1	BAL	A	14	1	-	0/1/2/3	0/0/0/0
1	AIB	A	16	1	-	0/2/3/6	0/0/0/0
1	BAL	A	17	1	-	0/1/2/3	0/0/0/0
1	AIB	A	19	1	-	0/2/3/6	0/0/0/0
1	AIB	A	2	1	-	0/2/3/6	0/0/0/0
1	AIB	A	20	1	-	0/2/3/6	0/0/0/0
1	BAL	A	22	1	-	0/1/1/3	0/0/0/0
1	AIB	A	4	1	-	0/2/3/6	0/0/0/0
1	DIV	A	7	1	-	0/3/6/9	0/0/0/0
1	AIB	B	10	1	-	0/2/3/6	0/0/0/0
1	BAL	B	11	1	-	0/1/2/3	0/0/0/0
1	DIV	B	13	1	-	0/3/6/9	0/0/0/0
1	BAL	B	14	1	-	0/1/2/3	0/0/0/0
1	AIB	B	16	1	-	0/2/3/6	0/0/0/0
1	BAL	B	17	1	-	0/1/2/3	0/0/0/0
1	AIB	B	19	1	-	0/2/3/6	0/0/0/0
1	AIB	B	2	1	-	0/2/3/6	0/0/0/0
1	AIB	B	20	1	-	0/2/3/6	0/0/0/0
1	BAL	B	22	1	-	0/1/2/3	0/0/0/0
1	AIB	B	4	1	-	0/2/3/6	0/0/0/0
1	DIV	B	7	1	-	0/3/6/9	0/0/0/0

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	AIB	O-C-CA	-4.00	118.07	124.62
1	B	4	AIB	O-C-CA	-3.76	118.47	124.62
1	B	10	AIB	O-C-CA	-3.51	118.88	124.62
1	A	10	AIB	O-C-CA	-3.38	119.10	124.62
1	B	4	AIB	CB2-CA-CB1	-3.18	105.54	110.42
1	B	20	AIB	O-C-CA	-3.05	119.63	124.62
1	A	4	AIB	O-C-CA	-2.84	119.97	124.62
1	B	2	AIB	CB2-CA-CB1	-2.75	106.20	110.42
1	A	7	DIV	O-C-CA	-2.70	118.41	125.01
1	B	19	AIB	O-C-CA	-2.53	120.48	124.62
1	A	13	DIV	O-C-CA	-2.28	119.44	125.01
1	B	2	AIB	O-C-CA	-2.23	120.97	124.62

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	7	DIV	O-C-CA	-2.23	119.57	125.01
1	A	16	AIB	O-C-CA	-2.22	120.99	124.62
1	A	20	AIB	O-C-CA	-2.17	121.08	124.62
1	B	13	DIV	O-C-CA	-2.13	119.80	125.01
1	A	19	AIB	O-C-CA	-2.09	121.21	124.62
1	A	10	AIB	CB2-CA-CB1	2.80	114.71	110.42

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	22	BAL	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	9/23 (39%)	28.63	9 (100%) 0 0	8, 10, 14, 14	0
1	B	10/23 (43%)	26.57	10 (100%) 0 0	9, 9, 12, 16	0
All	All	19/46 (41%)	27.55	19 (100%) 0 0	8, 9, 14, 16	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	LEU	36.2
1	A	6	PHE	33.2
1	B	18	LEU	29.6
1	A	5	PRO	29.5
1	B	21	LEU	29.4
1	A	21	LEU	29.1
1	A	15	LEU	28.9
1	B	6	PHE	28.9
1	B	15	LEU	28.4
1	A	9	ALA	28.2
1	B	9	ALA	27.9
1	B	23[A]	GLY	26.6
1	B	12	ALA	26.1
1	A	8	PRO	26.0
1	A	12	ALA	25.4
1	B	8	PRO	24.2
1	B	3	PRO	22.3
1	B	5	PRO	22.2
1	A	3	PRO	21.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	BAL	B	14	5/6	0.13	1.01	-	8,8,10,10	0
1	BAL	A	14	5/6	0.18	0.86	-	8,9,10,10	0
1	DIV	B	7	7/8	-0.01	1.01	-	8,11,18,18	0
1	AIB	B	2	6/7	0.10	0.97	-	10,17,25,25	0
1	AIB	B	10	6/7	-0.13	1.09	-	9,11,17,17	0
1	AIB	A	16	6/7	-0.05	1.02	-	7,9,13,13	0
1	AIB	B	16	6/7	-0.19	1.10	-	7,9,14,14	0
1	BAL	B	17	5/6	-0.01	1.05	-	8,9,10,10	0
1	BAL	A	22	4/6	-0.22	1.13	-	12,19,26,26	0
1	AIB	A	20	6/7	-0.06	1.13	-	9,11,16,16	0
1	AIB	B	19	6/7	0.15	0.91	-	9,11,18,18	0
1	BAL	B	11	5/6	-0.11	1.15	-	8,9,11,11	0
1	AIB	B	4	6/7	-0.24	1.27	-	8,11,17,17	0
1	DIV	A	13	7/8	-0.00	0.94	-	7,9,14,14	0
1	BAL	A	11	5/6	0.08	0.92	-	7,10,12,14	0
1	AIB	A	19	6/7	-0.01	1.05	-	8,10,16,16	0
1	AIB	A	10	6/7	0.01	1.00	-	12,17,27,27	0
1	DIV	B	13	7/8	-0.16	1.16	-	8,11,17,17	0
1	AIB	A	4	6/7	0.10	1.01	-	8,9,14,14	0
1	BAL	B	22	5/6	0.11	0.96	-	9,17,22,22	0
1	BAL	A	17	5/6	0.07	1.07	-	8,9,10,10	0
1	AIB	A	2	6/7	-0.12	1.07	-	10,14,22,22	0
1	DIV	A	7	7/8	-0.16	1.01	-	9,13,23,23	0
1	AIB	B	20	6/7	0.09	0.87	-	8,10,15,15	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.