



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:55 PM GMT

PDB ID : 1Z1A
Title : S. cerevisiae Sir1 ORC-interaction domain
Authors : Hou, Z.; Bernstein, D.A.; Fox, C.A.; Keck, J.L.
Deposited on : 2005-03-03
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	<div> <div>5%</div> <div>59%</div> <div>20%</div> <div>•</div> <div>18%</div> </div>
1	B	142	<div> <div>5%</div> <div>61%</div> <div>18%</div> <div>5%</div> <div>16%</div> </div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulatory protein SIR1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	117	Total	C	N	O	S	Se	0	1	0
			968	632	157	173	2	4			
1	B	119	Total	C	N	O	S	Se	0	0	0
			980	638	160	177	2	3			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	470	GLY	-	CLONING ARTIFACT	UNP P21691
A	471	SER	-	CLONING ARTIFACT	UNP P21691
A	472	HIS	-	CLONING ARTIFACT	UNP P21691
A	528	MSE	MET	MODIFIED RESIDUE	UNP P21691
A	532	MSE	MET	MODIFIED RESIDUE	UNP P21691
A	589	MSE	MET	MODIFIED RESIDUE	UNP P21691
B	470	GLY	-	CLONING ARTIFACT	UNP P21691
B	471	SER	-	CLONING ARTIFACT	UNP P21691
B	472	HIS	-	CLONING ARTIFACT	UNP P21691
B	528	MSE	MET	MODIFIED RESIDUE	UNP P21691
B	532	MSE	MET	MODIFIED RESIDUE	UNP P21691
B	589	MSE	MET	MODIFIED RESIDUE	UNP P21691
A	558	CYS	ALA	ENGINEERED	UNP P21691
B	558	CYS	ALA	ENGINEERED	UNP P21691


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	22	Total	O	0	0
			22	22		
2	B	21	Total	O	0	0
			21	21		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Regulatory protein SIR1



• Molecule 1: Regulatory protein SIR1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	60.93Å 60.93Å 159.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.50 24.19 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-2.50) 98.6 (24.19-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.16 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.238 , 0.274 0.240 , 0.268	Depositor DCC
R_{free} test set	1206 reflections (10.89%)	DCC
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.259	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 43.2	EDS
Estimated twinning fraction	0.035 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 12298 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1991	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/991	0.75	3/1333 (0.2%)
1	B	0.54	0/999	0.74	2/1345 (0.1%)
All	All	0.53	0/1990	0.74	5/2678 (0.2%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	498	ASP	CB-CG-OD2	6.67	124.30	118.30
1	B	498	ASP	CB-CG-OD2	5.72	123.44	118.30
1	A	610	ASP	CB-CG-OD2	5.13	122.92	118.30
1	B	610	ASP	CB-CG-OD2	5.11	122.90	118.30
1	A	539	ASP	CB-CG-OD2	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	968	0	997	21	0
1	B	980	0	1004	23	0
2	A	22	0	0	0	0
2	B	21	0	0	0	0
All	All	1991	0	2001	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:529:ILE:HA	1:A:532:MSE:HE2	1.50	0.94
1:B:529:ILE:HA	1:B:532:MSE:HE2	1.57	0.84
1:A:525:GLN:HA	1:A:528[A]:MSE:HE2	1.59	0.82
1:B:507:GLU:HB2	1:B:559:LYS:HE3	1.67	0.76
1:B:496:VAL:HG21	1:B:532:MSE:HE1	1.76	0.68
1:A:496:VAL:HG21	1:A:532:MSE:CE	2.24	0.67
1:A:501:LEU:HD23	1:A:510:ILE:HG23	1.76	0.67
1:B:496:VAL:HG21	1:B:532:MSE:CE	2.29	0.62
1:B:567:ARG:HH12	1:B:589:MSE:SE	2.31	0.62
1:B:567:ARG:NH1	1:B:589:MSE:SE	2.83	0.62
1:A:561:ILE:HG13	1:A:595:CYS:SG	2.43	0.57
1:A:541:LYS:NZ	1:A:601:ASP:HB2	2.21	0.55
1:A:496:VAL:HG21	1:A:532:MSE:HE3	1.90	0.54
1:B:606:GLN:HG3	1:B:607:TYR:H	1.73	0.54
1:A:485:THR:HG22	1:A:552:LYS:HB2	1.92	0.52
1:B:602:ASP:HB3	1:B:604:THR:HG23	1.94	0.50
1:B:487:GLU:HG2	1:B:553:THR:HG21	1.93	0.49
1:A:511:ASN:HD22	1:A:512:PRO:N	2.11	0.48
1:B:491:SER:HB2	1:B:492:PRO:CD	2.44	0.48
1:B:546:PRO:HA	1:B:606:GLN:HE21	1.80	0.47
1:A:532:MSE:HE3	1:A:534:LEU:HD21	1.96	0.47
1:B:606:GLN:HG3	1:B:607:TYR:N	2.29	0.47
1:A:496:VAL:HG21	1:A:532:MSE:HE1	1.96	0.47
1:B:546:PRO:HA	1:B:606:GLN:NE2	2.30	0.47
1:B:541:LYS:NZ	1:B:601:ASP:OD1	2.35	0.47
1:B:532:MSE:HE3	1:B:534:LEU:HD21	1.98	0.46
1:B:545:ASP:HB3	1:B:546:PRO:HD2	1.98	0.45
1:B:511:ASN:HD22	1:B:512:PRO:CD	2.29	0.45
1:B:491:SER:HB2	1:B:492:PRO:HD2	1.98	0.45
1:A:544:GLN:O	1:A:575:THR:OG1	2.34	0.44
1:A:507:GLU:HB2	1:A:559:LYS:HG3	2.00	0.43
1:B:572:LYS:NZ	1:B:611:ASP:OD2	2.51	0.43
1:B:529:ILE:HA	1:B:532:MSE:CE	2.40	0.43
1:B:511:ASN:HA	1:B:512:PRO:HD2	1.90	0.43
1:A:511:ASN:ND2	1:A:513:LYS:H	2.16	0.42
1:A:511:ASN:HD22	1:A:512:PRO:CD	2.32	0.42
1:A:563:LYS:HE2	1:A:607:TYR:CE1	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:524:HIS:O	1:B:528:MSE:HG2	2.20	0.42
1:A:541:LYS:HZ1	1:A:601:ASP:HB2	1.85	0.41
1:A:502:ILE:HD11	1:A:554:LEU:HD12	2.02	0.41
1:B:511:ASN:HD22	1:B:512:PRO:N	2.19	0.41
1:A:611:ASP:OD1	1:A:611:ASP:C	2.60	0.40
1:A:511:ASN:HD22	1:A:513:LYS:H	1.68	0.40
1:A:532:MSE:HE2	1:A:532:MSE:HB2	1.92	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/142 (80%)	110 (96%)	4 (4%)	0	100	100
1	B	115/142 (81%)	108 (94%)	7 (6%)	0	100	100
All	All	229/284 (81%)	218 (95%)	11 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/127 (87%)	102 (93%)	8 (7%)	17	32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	110/127 (87%)	99 (90%)	11 (10%)	9	18
All	All	220/254 (87%)	201 (91%)	19 (9%)	13	24

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	485	THR
1	A	495	LEU
1	A	511	ASN
1	A	532	MSE
1	A	536	LYS
1	A	538	ASN
1	A	575	THR
1	A	591	VAL
1	B	508	LYS
1	B	511	ASN
1	B	519	THR
1	B	528	MSE
1	B	538	ASN
1	B	559	LYS
1	B	567	ARG
1	B	579	GLU
1	B	589	MSE
1	B	601	ASP
1	B	602	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	511	ASN
1	A	524	HIS
1	A	538	ASN
1	B	511	ASN
1	B	606	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	114/142 (80%)	0.27	7 (6%) 25 27	29, 50, 87, 103	0
1	B	116/142 (81%)	0.24	7 (6%) 25 28	32, 54, 79, 88	0
All	All	230/284 (80%)	0.25	14 (6%) 25 27	29, 53, 82, 103	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	577	PRO	4.5
1	A	542	LYS	4.1
1	A	485	THR	3.3
1	A	595	CYS	3.2
1	B	490	VAL	3.1
1	B	533	ASN	2.8
1	A	484	SER	2.6
1	A	603	GLN	2.4
1	B	544	GLN	2.4
1	B	515	PRO	2.3
1	B	579	GLU	2.2
1	A	569	ALA	2.0
1	B	543	TYR	2.0
1	B	489	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.