



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:59 AM GMT

PDB ID : 2Z1U  
Title : Crystal Structure of Hydrogenase Maturation Protein HypE in complex with ATP  
Authors : Shomura, Y.; Higuchi, Y.  
Deposited on : 2007-05-15  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div>24%</div> <div>79%</div> <div>15%</div> <div>..</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2690 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase expression/formation protein HypE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2475	1542	443	475	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP Q72F88
A	0	ALA	-	EXPRESSION TAG	UNP Q72F88

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	182	Total	O	0	0
			182	182		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hydrogenase expression/formation protein HypE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.63Å 67.63Å 184.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.30 – 2.00 42.38 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (33.30-2.00) 99.2 (42.38-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.66 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.200 , 0.228 0.199 , 0.223	Depositor DCC
$R_{free}$ test set	1707 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.1	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.8	EDS
Estimated twinning fraction	0.030 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 33651 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2690	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/2510	0.73	3/3399 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	221	ARG	NE-CZ-NH2	-5.87	117.37	120.30
1	A	221	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	327	ARG	NE-CZ-NH2	-5.17	117.72	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	226	THR	Mainchain
1	A	46	THR	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2475	0	2497	44	0
2	A	2	0	0	0	0
3	A	31	0	12	1	0
4	A	182	0	0	3	0
All	All	2690	0	2509	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:VAL:HG11	1:A:139:MET:CE	2.03	0.89
1:A:221:ARG:HD3	3:A:342:ATP:O2G	1.76	0.86
1:A:139:MET:HE2	1:A:141:ILE:HD11	1.57	0.86
1:A:226:THR:HG23	1:A:331:MET:HG2	1.61	0.82
1:A:36:THR:HG21	4:A:357:HOH:O	1.81	0.80
1:A:108:VAL:HG11	1:A:139:MET:HE2	1.63	0.80
1:A:219:PRO:HA	1:A:227:THR:HG21	1.66	0.77
1:A:139:MET:CE	1:A:141:ILE:HD11	2.15	0.77
1:A:178:THR:HG21	1:A:193:CYS:C	2.07	0.74
1:A:108:VAL:HG11	1:A:139:MET:HE1	1.72	0.71
1:A:227:THR:HG22	4:A:513:HOH:O	1.91	0.69
1:A:219:PRO:HA	1:A:227:THR:CG2	2.28	0.62
1:A:321:THR:HG22	1:A:323:LEU:H	1.65	0.62
1:A:217:ARG:HH22	1:A:234:GLN:HE22	1.47	0.60
1:A:174:ASP:O	1:A:178:THR:CG2	2.51	0.58
1:A:70:THR:HG23	1:A:196:SER:OG	2.04	0.58
1:A:226:THR:HG23	1:A:331:MET:CG	2.32	0.58
1:A:178:THR:HG21	1:A:193:CYS:CA	2.34	0.57
1:A:174:ASP:O	1:A:178:THR:HG22	2.05	0.57
1:A:138:LYS:HE2	4:A:419:HOH:O	2.05	0.56
1:A:5:THR:HB	1:A:102:MET:HG2	1.90	0.54
1:A:321:THR:CG2	1:A:323:LEU:H	2.22	0.52
1:A:220:THR:O	1:A:267:ALA:HA	2.10	0.51
1:A:310:GLY:O	1:A:311:ALA:HB3	2.12	0.48
1:A:227:THR:OG1	1:A:273:ILE:HD13	2.13	0.48
1:A:310:GLY:O	1:A:311:ALA:CB	2.62	0.47
1:A:114:LYS:O	1:A:118:GLU:HG3	2.16	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:HIS:CE1	1:A:309:LEU:HD12	2.51	0.45
1:A:178:THR:HG21	1:A:193:CYS:HA	1.98	0.45
1:A:157:GLY:HA3	1:A:234:GLN:NE2	2.32	0.44
1:A:217:ARG:HH12	1:A:234:GLN:NE2	2.16	0.44
1:A:226:THR:O	1:A:230:GLU:HG3	2.19	0.43
1:A:212:ASP:HB3	1:A:280:ARG:HE	1.83	0.43
1:A:174:ASP:OD1	1:A:249:ARG:HD2	2.19	0.42
1:A:238:VAL:HG22	1:A:240:HIS:CD2	2.54	0.42
1:A:14:GLY:O	1:A:18:HIS:ND1	2.46	0.42
1:A:174:ASP:O	1:A:178:THR:HG23	2.19	0.42
1:A:321:THR:HG22	1:A:323:LEU:N	2.34	0.42
1:A:156:SER:HB3	1:A:159:ARG:HG3	2.00	0.42
1:A:61:ILE:HG23	1:A:68:ILE:HG22	2.01	0.42
1:A:177:LEU:HD12	1:A:252:VAL:HA	2.03	0.41
1:A:218:ASP:N	1:A:218:ASP:OD1	2.52	0.41
1:A:56:TYR:CE1	1:A:72:ALA:HA	2.56	0.41
1:A:226:THR:CG2	1:A:331:MET:HG2	2.41	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/343 (96%)	316 (96%)	12 (4%)	3 (1%)	21	13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	332	LEU
1	A	30	ASP
1	A	311	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	264/268 (98%)	241 (91%)	23 (9%)	13 7

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	9	ASP
1	A	20	LEU
1	A	36	THR
1	A	43	LEU
1	A	70	THR
1	A	71	LEU
1	A	90	LEU
1	A	100	LEU
1	A	121	VAL
1	A	177	LEU
1	A	178	THR
1	A	179	ILE
1	A	180	LEU
1	A	207	VAL
1	A	227	THR
1	A	238	VAL
1	A	280	ARG
1	A	285	LEU
1	A	313	ARG
1	A	318	VAL
1	A	321	THR
1	A	327	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	229	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	234	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ATP	A	342	2	24,33,33	1.03	2 (8%)	31,52,52	2.38	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	342	2	-	0/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	342	ATP	C2-N3	2.11	1.35	1.32
3	A	342	ATP	O4'-C1'	2.74	1.44	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	342	ATP	N3-C2-N1	-10.77	120.65	128.89
3	A	342	ATP	PA-O3A-PB	-3.60	122.62	132.73
3	A	342	ATP	C1'-N9-C4	-2.71	122.85	126.94
3	A	342	ATP	C2-N1-C6	3.07	124.25	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	342	ATP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/343 (97%)	1.33	82 (24%) <b>1</b> <b>1</b>	44, 52, 65, 81	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	PHE	6.6
1	A	143	THR	6.3
1	A	56	TYR	6.1
1	A	141	ILE	6.0
1	A	90	LEU	5.7
1	A	161	ARG	5.6
1	A	93	GLY	5.6
1	A	55	SER	5.5
1	A	309	LEU	5.5
1	A	144	THR	5.3
1	A	94	PHE	5.2
1	A	95	ILE	5.0
1	A	91	SER	5.0
1	A	190	ALA	5.0
1	A	53	THR	5.0
1	A	92	CYS	4.9
1	A	313	ARG	4.8
1	A	333	GLU	4.7
1	A	72	ALA	4.7
1	A	142	ASN	4.5
1	A	77	VAL	4.4
1	A	187	ASN	4.2
1	A	76	THR	4.1
1	A	186	LEU	4.0
1	A	145	GLY	4.0
1	A	73	VAL	4.0
1	A	312	ALA	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	189	ALA	3.7
1	A	68	ILE	3.7
1	A	71	LEU	3.7
1	A	306	VAL	3.7
1	A	54	ASP	3.7
1	A	305	SER	3.7
1	A	125	THR	3.7
1	A	47	GLY	3.6
1	A	126	GLY	3.6
1	A	146	ILE	3.6
1	A	307	GLY	3.5
1	A	332	LEU	3.5
1	A	123	ILE	3.4
1	A	308	GLU	3.4
1	A	226	THR	3.4
1	A	75	GLY	3.3
1	A	80	VAL	3.3
1	A	51	MET	3.2
1	A	127	ASP	3.2
1	A	212	ASP	3.2
1	A	121	VAL	3.2
1	A	7	LEU	3.2
1	A	133	ARG	3.1
1	A	58	VAL	3.1
1	A	128	THR	3.1
1	A	129	LYS	3.0
1	A	314	ALA	3.0
1	A	258	PHE	3.0
1	A	57	THR	2.9
1	A	160	ALA	2.9
1	A	96	LEU	2.8
1	A	140	PHE	2.8
1	A	341	CYS	2.7
1	A	112	MET	2.7
1	A	103	ASP	2.6
1	A	78	ASN	2.6
1	A	311	ALA	2.6
1	A	152	ASP	2.4
1	A	310	GLY	2.4
1	A	185	GLY	2.3
1	A	272	LEU	2.3
1	A	52	SER	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	250	GLU	2.3
1	A	304	LYS	2.2
1	A	199	LEU	2.2
1	A	124	VAL	2.2
1	A	236	GLN	2.2
1	A	220	THR	2.1
1	A	40	ALA	2.1
1	A	153	PRO	2.1
1	A	74	HIS	2.1
1	A	240	HIS	2.1
1	A	340	ILE	2.1
1	A	41	ALA	2.1
1	A	69	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	344	1/1	0.81	0.19	-0.77	47,47,47,47	0
3	ATP	A	342	31/31	0.96	0.10	-1.87	24,31,34,35	0
2	MG	A	343	1/1	0.97	0.09	-3.74	29,29,29,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.