



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:00 AM GMT

PDB ID : 2Z63
Title : Crystal structure of the TV8 hybrid of human TLR4 and hagfish VLRB.61
Authors : Kim, H.M.; Park, B.S.; Lee, J.-O.
Deposited on : 2007-07-22
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	570	<div> <div>4%</div> <div>78%</div> <div>20%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	1412	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 4, Variable lymphocyte receptor B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	570	Total	C	N	O	S	0	0	0
			4536	2901	754	858	23			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	528	ALA	-	SEE REMARK 999	UNP O00206
A	529	SER	-	SEE REMARK 999	UNP O00206

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	3	Total	C	N	O	0	0
			38	22	2	14		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	292	Total	O	0	0
			292	292		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Toll-like receptor 4, Variable lymphocyte receptor B



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	88.14Å 88.14Å 92.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.00 38.16 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.8 (20.00-2.00) 95.8 (38.16-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.250 , 0.298 0.251 , 0.292	Depositor DCC
R_{free} test set	2622 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 29.6	EDS
Estimated twinning fraction	0.237 for -h,-k,l 0.387 for h,-h-k,-l 0.218 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.32$, $\langle L^2 \rangle = 0.15$	Xtriage
Outliers	0 of 51622 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4894	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/4631	0.66	1/6269 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	283	LEU	CA-CB-CG	5.84	128.73	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4536	0	4504	72	0
2	A	38	0	34	1	0
3	A	28	0	25	1	0
4	A	292	0	0	9	0
All	All	4894	0	4563	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:378:LEU:HD22	1:A:404:LEU:HD12	1.55	0.89
1:A:418:PHE:HB3	4:A:1187:HOH:O	1.76	0.82
1:A:235:ASN:HD21	1:A:263:PHE:H	1.32	0.77
1:A:378:LEU:HD22	1:A:404:LEU:CD1	2.13	0.76
1:A:479:ALA:HB1	4:A:1041:HOH:O	1.85	0.75
1:A:269:LEU:HD12	1:A:269:LEU:H	1.53	0.74
1:A:221:GLY:HA2	1:A:224:LYS:HE3	1.71	0.73
1:A:77:PHE:HB3	1:A:80:LEU:HG	1.70	0.73
1:A:421:LEU:HB2	4:A:1187:HOH:O	1.91	0.70
1:A:388:LYS:HD2	1:A:388:LYS:H	1.59	0.68
1:A:195:LEU:HD11	1:A:222:ALA:O	1.94	0.67
1:A:229:HIS:O	1:A:256:HIS:HB3	1.95	0.65
3:A:1411:NAG:H3	3:A:1411:NAG:H83	1.78	0.65
1:A:221:GLY:CA	1:A:224:LYS:HE3	2.27	0.64
1:A:159:HIS:HA	1:A:184:SER:O	1.97	0.64
1:A:317:SER:HA	1:A:339:ASN:O	1.97	0.64
1:A:516:PHE:HB3	1:A:543:LEU:HD21	1.78	0.63
1:A:154:GLU:HG3	1:A:179:HIS:HB3	1.83	0.60
1:A:585:GLY:HA2	4:A:1227:HOH:O	2.02	0.59
1:A:320:ILE:HG23	1:A:323:VAL:HG23	1.84	0.58
1:A:197:VAL:HG13	4:A:1092:HOH:O	2.02	0.58
2:A:1402:FUL:H3	2:A:1403:NAG:H82	1.87	0.57
1:A:447:ARG:NH2	4:A:1031:HOH:O	2.38	0.57
1:A:233:LEU:HA	1:A:236:ASN:HD21	1.70	0.56
1:A:383:ASN:HB2	1:A:385:LEU:HG	1.88	0.55
1:A:297:LEU:HD13	1:A:303:LEU:HD13	1.88	0.55
1:A:235:ASN:ND2	1:A:263:PHE:H	2.02	0.55
1:A:344:GLN:HA	1:A:364:GLY:HA3	1.89	0.54
1:A:316:VAL:HG22	1:A:338:VAL:HG22	1.90	0.53
1:A:393:GLN:HG3	1:A:420:GLY:HA3	1.89	0.53
1:A:391:CYS:O	1:A:418:PHE:HA	2.09	0.53
1:A:148:HIS:H	1:A:148:HIS:CD2	2.27	0.53
1:A:67:ARG:HA	1:A:89:GLU:O	2.09	0.52
1:A:67:ARG:HB3	1:A:68:HIS:HD2	1.75	0.52
1:A:436:GLN:NE2	1:A:436:GLN:HA	2.24	0.51
1:A:44:ASN:ND2	4:A:1116:HOH:O	2.44	0.51
1:A:80:LEU:HB3	1:A:104:LEU:CD1	2.41	0.51
1:A:386:SER:HB2	1:A:388:LYS:NZ	2.26	0.51
1:A:93:ILE:HG22	1:A:121:ALA:HB1	1.93	0.50
1:A:179:HIS:ND1	1:A:207:SER:HB2	2.26	0.50
1:A:61:LEU:O	1:A:88:CYS:SG	2.71	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:LEU:HB3	1:A:283:LEU:HD11	1.95	0.48
1:A:98:TYR:HB3	1:A:101:LEU:HD12	1.94	0.47
1:A:156:ASN:HA	1:A:181:ASP:HB3	1.95	0.47
1:A:254:GLU:HA	1:A:284:THR:HB	1.97	0.47
1:A:125:LEU:HD13	1:A:128:LEU:HD22	1.96	0.47
1:A:386:SER:HB2	1:A:388:LYS:HZ1	1.80	0.46
1:A:227:ARG:HA	1:A:254:GLU:O	2.15	0.46
1:A:524:VAL:HG13	1:A:550:TRP:HZ3	1.79	0.46
1:A:114:ILE:HD12	1:A:136:THR:HB	1.99	0.45
1:A:505:GLN:HB2	4:A:1041:HOH:O	2.15	0.45
1:A:422:GLU:CD	1:A:422:GLU:H	2.20	0.45
1:A:342:PHE:C	1:A:344:GLN:H	2.20	0.45
1:A:287:GLU:HG2	1:A:312:SER:HB3	1.97	0.45
1:A:393:GLN:CG	1:A:420:GLY:HA3	2.46	0.44
1:A:232:THR:HG22	1:A:234:ARG:HG3	2.00	0.44
1:A:334:HIS:HE1	1:A:336:GLU:HB2	1.82	0.44
1:A:110:THR:HG23	1:A:134:VAL:HB	1.99	0.43
1:A:310:VAL:HG12	1:A:332:TRP:CD1	2.53	0.43
1:A:143:ASN:O	1:A:145:PRO:HD3	2.19	0.43
1:A:185:ASN:HB3	1:A:186:LYS:H	1.67	0.43
1:A:205:ASN:OD1	1:A:229:HIS:HB2	2.18	0.43
1:A:234:ARG:HA	1:A:261:GLY:O	2.19	0.43
1:A:306:CYS:HB2	4:A:1150:HOH:O	2.19	0.42
1:A:390:CYS:HA	1:A:391:CYS:HA	1.90	0.42
1:A:313:PHE:HE2	1:A:315:LEU:HG	1.85	0.42
1:A:313:PHE:CE2	1:A:315:LEU:HG	2.55	0.41
1:A:158:ALA:HA	1:A:183:SER:O	2.20	0.41
1:A:120:GLY:O	1:A:123:SER:HB2	2.20	0.41
1:A:48:ILE:H	1:A:48:ILE:HD12	1.85	0.41
1:A:436:GLN:NE2	1:A:440:PHE:HE1	2.18	0.41
1:A:412:ILE:HB	1:A:434:LEU:HD23	2.02	0.41
1:A:233:LEU:CA	1:A:236:ASN:ND2	2.84	0.40
1:A:95:ASP:HA	1:A:120:GLY:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	568/570 (100%)	520 (92%)	41 (7%)	7 (1%)	16	8

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	89	GLU
1	A	276	ALA
1	A	35	ASN
1	A	42	GLU
1	A	65	PRO
1	A	267	GLY
1	A	364	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	526/526 (100%)	498 (95%)	28 (5%)	28	22

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LEU
1	A	95	ASP
1	A	105	SER
1	A	115	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	123	SER
1	A	148	HIS
1	A	156	ASN
1	A	195	LEU
1	A	216	ASN
1	A	250	LEU
1	A	255	VAL
1	A	266	GLU
1	A	269	LEU
1	A	277	LEU
1	A	278	GLU
1	A	293	LEU
1	A	301	ILE
1	A	307	LEU
1	A	308	THR
1	A	324	LYS
1	A	329	ASN
1	A	374	SER
1	A	388	LYS
1	A	393	GLN
1	A	402	LYS
1	A	446	LEU
1	A	485	GLU
1	A	575	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	156	ASN
1	A	216	ASN
1	A	235	ASN
1	A	236	ASN
1	A	268	ASN
1	A	329	ASN
1	A	393	GLN
1	A	426	HIS
1	A	436	GLN
1	A	484	GLN
1	A	510	GLN
1	A	523	GLN
1	A	547	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	578	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	1401	1,2	14,14,15	0.59	0	15,19,21	1.06	1 (6%)
2	FUL	A	1402	2	10,10,11	0.59	0	14,14,16	1.04	1 (7%)
2	NAG	A	1403	2	14,14,15	0.57	0	15,19,21	0.93	1 (6%)
3	NAG	A	1411	1,3	14,14,15	0.68	0	15,19,21	1.38	1 (6%)
3	NAG	A	1412	3	14,14,15	0.54	0	15,19,21	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1401	1,2	-	0/6/23/26	0/1/1/1
2	FUL	A	1402	2	-	0/0/17/20	0/1/1/1
2	NAG	A	1403	2	-	0/6/23/26	0/1/1/1
3	NAG	A	1411	1,3	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1412	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1401	NAG	C1-O5-C5	2.16	114.98	112.25
2	A	1403	NAG	C1-O5-C5	2.16	114.99	112.25
2	A	1402	FUL	O5-C5-C6	3.20	111.43	106.13
3	A	1411	NAG	C2-N2-C7	3.35	127.34	123.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1402	FUL	1	0
2	A	1403	NAG	1	0
3	A	1411	NAG	1	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	570/570 (100%)	0.51	23 (4%)	42 44	21, 31, 40, 51	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	GLU	5.3
1	A	202	PRO	4.7
1	A	322	ARG	4.5
1	A	103	HIS	4.4
1	A	240	LEU	3.6
1	A	364	GLY	3.5
1	A	575	GLN	3.5
1	A	198	LEU	3.2
1	A	301	ILE	3.2
1	A	326	PHE	3.0
1	A	29	CYS	2.9
1	A	46	TYR	2.7
1	A	199	HIS	2.6
1	A	30	VAL	2.6
1	A	328	TYR	2.6
1	A	583	CYS	2.6
1	A	323	VAL	2.4
1	A	365	ASN	2.4
1	A	50	ASP	2.4
1	A	68	HIS	2.1
1	A	278	GLU	2.1
1	A	53	PRO	2.1
1	A	321	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	A	1412	14/15	0.79	0.22	3.19	39,47,56,58	0
3	NAG	A	1411	14/15	0.94	0.11	-	16,26,33,39	0
2	FUL	A	1402	10/11	0.56	0.72	-	64,66,68,68	0
2	NAG	A	1401	14/15	0.74	0.27	-	42,50,62,66	0
2	NAG	A	1403	14/15	0.46	0.43	-	67,70,74,76	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.