



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 4, 2016 – 01:41 PM EDT

PDB ID : 4Z7M  
Title : Novel Inhibitors of Bacterial Methionine Aminopeptidase with Broad-Spectrum Biochemical Activity  
Authors : Rose, J.A.; Lahiri, S.D.; McKinney, D.C.; Albert, R.; Morningstar, M.; Shapiro, A.B.; Fisher, S.F.; Fleming, P.R.  
Deposited on : 2015-04-07  
Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1164 (1.46-1.42)
Clashscore	102246	1219 (1.46-1.42)
Ramachandran outliers	100387	1200 (1.46-1.42)
Sidechain outliers	100360	1200 (1.46-1.42)
RSRZ outliers	91569	1166 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	
1	B	261	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Methionine aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	0	0
			2026	1277	350	384	15			
1	B	261	Total	C	N	O	S	0	0	0
			2026	1277	350	384	15			

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is N 2 -[(3,5-difluorophenyl)acetyl]-N-[(3S,7R)-1-methyl-2-oxo-7-phenyl-2,3,4,7-tetrahydro-1H-azepin-3-yl]-L-alaninamide (three-letter code: 4L9) (formula: C<sub>24</sub>H<sub>25</sub>F<sub>2</sub>N<sub>3</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			32	24	2	3	3		
3	B	1	Total	C	F	N	O	0	0
			32	24	2	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	192	Total	O	0	0
			192	192		
4	B	195	Total	O	0	0
			195	195		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methionine aminopeptidase

Chain A:  88% 9% .



- Molecule 1: Methionine aminopeptidase

Chain B:  89% 9% .



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.98Å 62.71Å 72.98Å 90.00° 89.59° 90.00°	Depositor
Resolution (Å)	47.57 – 1.43 47.56 – 1.43	Depositor EDS
% Data completeness (in resolution range)	69.6 (47.57-1.43) 68.8 (47.56-1.43)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 1.43Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.251 , 0.305 0.259 , 0.311	Depositor DCC
$R_{free}$ test set	3589 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.1	Xtriage
Anisotropy	0.505	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 43.0	EDS
Estimated twinning fraction	0.000 for k,h,-l 0.000 for -k,-h,-l 0.389 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 71301 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4505	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4L9, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	2/2061 (0.1%)	0.98	8/2785 (0.3%)
1	B	0.87	1/2061 (0.0%)	1.01	10/2785 (0.4%)
All	All	0.85	3/4122 (0.1%)	0.99	18/5570 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	204	GLU	CD-OE1	6.78	1.33	1.25
1	B	204	GLU	CD-OE1	6.37	1.32	1.25
1	A	235	GLU	CD-OE2	-5.13	1.20	1.25

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	15	ARG	NE-CZ-NH2	-9.91	115.34	120.30
1	B	15	ARG	NE-CZ-NH1	9.68	125.14	120.30
1	A	166	ARG	NE-CZ-NH1	7.72	124.16	120.30
1	B	228	ARG	NE-CZ-NH2	7.51	124.05	120.30
1	A	124	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	A	124	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	A	219	ASP	CB-CG-OD1	5.67	123.40	118.30
1	B	138	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	43	ARG	NE-CZ-NH2	-5.59	117.51	120.30
1	B	177	PHE	CB-CG-CD1	5.57	124.70	120.80
1	A	15	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	A	228	ARG	NE-CZ-NH1	-5.45	117.57	120.30
1	A	15	ARG	NE-CZ-NH1	5.39	122.99	120.30
1	B	83	ASP	CB-CG-OD1	5.34	123.11	118.30
1	B	19	ARG	NE-CZ-NH1	5.21	122.91	120.30
1	B	27	MET	CG-SD-CE	-5.19	91.90	100.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	97	ASP	CB-CG-OD1	5.15	122.94	118.30
1	B	43	ARG	NE-CZ-NH2	-5.05	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2026	0	2049	26	0
1	B	2026	0	2049	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	32	0	0	0	0
3	B	32	0	0	0	0
4	A	192	0	0	13	0
4	B	195	0	0	5	0
All	All	4505	0	4098	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:ILE:HD12	4:A:558:HOH:O	1.31	1.23
1:A:112:MET:CE	4:A:558:HOH:O	2.13	0.97
1:A:93:ILE:CG1	4:A:558:HOH:O	2.18	0.90
1:A:93:ILE:CD1	4:A:558:HOH:O	1.97	0.85
1:A:112:MET:HE2	4:A:558:HOH:O	1.76	0.82
1:A:166:ARG:HG2	1:A:166:ARG:HH11	1.55	0.72
1:A:216:THR:HG23	4:A:462:HOH:O	1.91	0.70
1:B:190:GLU:OE2	4:B:401:HOH:O	2.10	0.69
1:A:216:THR:CG2	4:A:462:HOH:O	2.42	0.66
1:B:124:ARG:NH2	1:B:160:GLU:OE1	2.27	0.66

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:GLU:OE1	1:A:127:ARG:NH1	2.31	0.63
1:B:138:ARG:HD2	4:B:557:HOH:O	1.99	0.61
1:B:74:ASN:O	4:B:402:HOH:O	2.16	0.60
1:B:2:ALA:HB3	1:B:180:GLU:OE2	2.07	0.54
1:A:247:ILE:HG21	1:A:250:LEU:HD13	1.90	0.53
1:A:124:ARG:NH2	1:A:160:GLU:OE1	2.41	0.52
1:A:112:MET:HE1	4:A:558:HOH:O	1.95	0.52
1:B:169:CYS:HB2	1:B:183:VAL:O	2.11	0.50
1:B:3:ILE:HD11	1:B:181:PRO:HB3	1.93	0.50
1:A:163:SER:OG	1:A:208:ASN:HB2	2.14	0.47
1:A:93:ILE:HB	4:A:558:HOH:O	2.14	0.47
1:A:93:ILE:CB	4:A:558:HOH:O	2.60	0.47
1:B:262:HIS:HE1	4:B:415:HOH:O	1.98	0.47
1:A:166:ARG:HH11	1:A:166:ARG:CG	2.27	0.46
1:A:185:HIS:CE1	1:A:204:GLU:H	2.34	0.46
1:B:33:LYS:HB3	4:B:520:HOH:O	2.16	0.46
1:A:93:ILE:HG13	4:A:558:HOH:O	2.04	0.45
1:A:9:GLU:HB2	4:A:484:HOH:O	2.17	0.44
1:A:51:ASN:OD1	1:B:86:LYS:NZ	2.50	0.43
1:A:247:ILE:CG2	1:A:250:LEU:HD13	2.48	0.43
1:A:262:HIS:HE1	4:A:448:HOH:O	2.02	0.43
1:B:15:ARG:HD3	1:B:246:GLU:OE2	2.19	0.42
1:A:185:HIS:CD2	1:A:205:PRO:HA	2.55	0.42
1:B:35:GLY:O	1:B:87:LEU:HD22	2.21	0.41
1:A:214:ILE:C	1:A:214:ILE:HD12	2.41	0.41
1:B:75:GLU:CD	1:B:75:GLU:H	2.23	0.41
1:A:140:VAL:HG11	1:A:238:ILE:HD12	2.02	0.40
1:A:250:LEU:HD22	1:A:258:ALA:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/261 (99%)	251 (97%)	8 (3%)	0	100	100
1	B	259/261 (99%)	252 (97%)	7 (3%)	0	100	100
All	All	518/522 (99%)	503 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/225 (100%)	216 (96%)	9 (4%)	38	6
1	B	225/225 (100%)	222 (99%)	3 (1%)	76	44
All	All	450/450 (100%)	438 (97%)	12 (3%)	52	15

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLU
1	A	166	ARG
1	A	177	PHE
1	A	189	ARG
1	A	190	GLU
1	A	216	THR
1	A	228	ARG
1	A	243	ASN
1	A	250	LEU
1	B	27	MET
1	B	56	VAL
1	B	177	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	185	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	243	ASN
1	B	51	ASN
1	B	95	ASN
1	B	185	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	4L9	A	302	-	32,34,34	1.57	6 (18%)	36,47,47	2.42	8 (22%)
3	4L9	B	302	-	32,34,34	1.54	5 (15%)	36,47,47	2.44	9 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4L9	A	302	-	-	0/20/38/38	0/3/3/3
3	4L9	B	302	-	-	0/20/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	4L9	C15-N14	-4.37	1.30	1.35
3	A	302	4L9	C15-N14	-4.34	1.30	1.35
3	A	302	4L9	C11-C10	-4.04	1.22	1.33
3	B	302	4L9	C11-C10	-3.23	1.24	1.33
3	A	302	4L9	C7-C15	-2.35	1.50	1.53
3	A	302	4L9	C29-C30	2.05	1.41	1.37
3	A	302	4L9	C18-C12	2.27	1.55	1.52
3	B	302	4L9	C18-C12	2.28	1.55	1.52
3	B	302	4L9	C29-C30	2.40	1.41	1.37
3	A	302	4L9	C9-C7	2.80	1.57	1.53
3	B	302	4L9	C9-C7	3.43	1.58	1.53

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	4L9	O16-C15-N14	-6.70	113.17	121.65
3	B	302	4L9	O16-C15-N14	-4.77	115.62	121.65
3	A	302	4L9	C27-C28-C33	-2.85	115.56	120.37
3	B	302	4L9	C2-N24-C25	-2.74	117.19	121.42
3	B	302	4L9	O16-C15-C7	-2.58	115.12	119.88
3	B	302	4L9	C1-C2-N24	-2.54	105.44	110.27
3	A	302	4L9	C19-C18-C12	-2.06	117.09	120.70
3	B	302	4L9	C27-C28-C33	-2.01	116.98	120.37
3	A	302	4L9	F34-C32-C31	2.11	121.06	118.22
3	A	302	4L9	C12-C11-C10	2.38	131.32	123.17
3	B	302	4L9	C27-C28-C29	2.59	124.75	120.37
3	A	302	4L9	C27-C28-C29	2.61	124.79	120.37
3	A	302	4L9	C23-C18-C12	2.79	125.59	120.70
3	B	302	4L9	C12-C11-C10	3.06	133.67	123.17
3	B	302	4L9	C28-C33-C32	3.29	121.63	118.88
3	A	302	4L9	C7-C15-N14	10.62	128.99	117.75
3	B	302	4L9	C7-C15-N14	10.85	129.23	117.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	261/261 (100%)	-0.19	1 (0%) 93 95	23, 31, 42, 69	0
1	B	261/261 (100%)	-0.26	0 100 100	21, 30, 41, 61	0
All	All	522/522 (100%)	-0.23	1 (0%) 95 96	21, 30, 42, 69	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	ARG	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	4L9	A	302	32/32	0.90	0.11	1.88	23,28,36,46	0
3	4L9	B	302	32/32	0.90	0.09	1.11	23,26,34,37	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MN	B	301	1/1	1.00	0.07	-1.12	25,25,25,25	0
2	MN	A	301	1/1	0.98	0.03	-4.82	28,28,28,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.