Full wwPDB X-ray Structure Validation Report (i)

May 4, 2016 – 05:30 PM EDT

PDB ID	:	4ZB5
Title	:	A form of glucose isomerase collected at 100K.
Authors	:	Sandy, J.
Deposited on	:	2015-04-14
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027457
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3649 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylose isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	387	3123	1964	571	580	8	0	10	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	388	GLY	-	expression tag	UNP P24300

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
2	A	2	2	2	0	0

- Molecule 3 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	12	6	6	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	509	512	512	0	3

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylose isomerase

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α, β, γ	92.90Å 98.15Å 102.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.90 – 2.00 68.90 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (68.90-2.00) 99.7 (68.90-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	14.27 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.115 , 0.154 0.115 , 0.155	Depositor DCC
R_{free} test set	1540 reflections (5.06%)	DCC
Wilson B-factor (Å^2)	8.4	Xtriage
Anisotropy	0.360	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 53.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -h,-l,-k	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	3649	wwPDB-VP
Average B, all atoms (Å^2)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.54% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	2.17	22/3223 (0.7%)	1.38	39/4356 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	74[A]	ARG	NE-CZ	72.69	2.27	1.33
1	A	74[B]	ARG	NE-CZ	72.69	2.27	1.33
1	A	160	GLU	CD-OE1	10.47	1.37	1.25
1	A	328[A]	GLU	CD-OE2	9.78	1.36	1.25
1	A	328[B]	GLU	CD-OE2	9.78	1.36	1.25
1	A	160	GLU	CD-OE2	6.79	1.33	1.25
1	A	207	GLU	CD-OE2	6.79	1.33	1.25
1	A	340	ARG	CZ-NH2	6.70	1.41	1.33
1	A	74[A]	ARG	CZ-NH1	6.63	1.41	1.33
1	A	74[B]	ARG	CZ-NH1	6.63	1.41	1.33
1	A	337	GLU	CD-OE2	-6.04	1.19	1.25
1	A	128	GLU	CD-OE1	5.85	1.32	1.25
1	A	128	GLU	CD-OE2	5.71	1.31	1.25
1	A	266	ARG	CD-NE	-5.63	1.36	1.46
1	A	8	GLU	CG-CD	5.48	1.60	1.51
1	A	325	GLU	CG-CD	5.45	1.60	1.51
1	A	315	GLU	CD-OE2	-5.40	1.19	1.25
1	A	128	GLU	CG-CD	5.37	1.60	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	217	GLU	CD-OE1	-5.28	1.19	1.25
1	A	74[A]	ARG	CZ-NH2	5.25	1.39	1.33
1	A	74[B]	ARG	CZ-NH2	5.25	1.39	1.33
1	A	20	TRP	CB-CG	5.06	1.59	1.50

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74[A]	ARG	CD-NE-CZ	-25.41	88.03	123.60
1	A	74[B]	ARG	CD-NE-CZ	-25.41	88.03	123.60
1	A	74[A]	ARG	NE-CZ-NH2	19.87	130.24	120.30
1	A	74[B]	ARG	NE-CZ-NH2	19.87	130.24	120.30
1	A	340	ARG	NE-CZ-NH1	-11.94	114.33	120.30
1	A	340	ARG	NE-CZ-NH2	11.52	126.06	120.30
1	A	266	ARG	NE-CZ-NH2	-10.34	115.13	120.30
1	A	74[A]	ARG	NH1-CZ-NH2	-9.15	109.33	119.40
1	A	74[B]	ARG	NH1-CZ-NH2	-9.15	109.33	119.40
1	A	354	ARG	NE-CZ-NH1	8.67	124.63	120.30
1	A	266	ARG	NE-CZ-NH1	8.18	124.39	120.30
1	A	368[A]	ARG	NE-CZ-NH2	-7.21	116.69	120.30
1	A	368[B]	ARG	NE-CZ-NH2	-7.21	116.69	120.30
1	A	295	ASP	CB-CG-OD1	6.96	124.56	118.30
1	A	113	ARG	NE-CZ-NH2	-6.93	116.83	120.30
1	A	245	ASP	CB-CG-OD1	6.64	124.28	118.30
1	A	113	ARG	NE-CZ-NH1	6.37	123.48	120.30
1	A	337	GLU	CG-CD-OE1	6.36	131.01	118.30
1	A	160	GLU	OE1-CD-OE2	6.32	130.89	123.30
1	A	336	ASP	CB-CG-OD1	6.28	123.95	118.30
1	A	337	GLU	CG-CD-OE2	-6.15	106.01	118.30
1	A	24	ASP	CB-CG-OD2	-6.05	112.86	118.30
1	A	295	ASP	CB-CG-OD2	-5.78	113.10	118.30
1	A	368[A]	ARG	NE-CZ-NH1	5.77	123.19	120.30
1	A	368[B]	ARG	NE-CZ-NH1	5.77	123.19	120.30
1	A	381	ASP	CB-CG-OD1	5.60	123.34	118.30
1	A	35	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	124	ASP	CB-CG-OD1	5.48	123.23	118.30
1	A	150	ASP	CB-CG-OD1	5.32	123.09	118.30
1	A	74[A]	ARG	NE-CZ-NH1	-5.32	117.64	120.30
1	A	74[B]	ARG	NE-CZ-NH1	-5.32	117.64	120.30
1	A	157	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	A	112	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	188	ARG	NE-CZ-NH2	-5.21	117.69	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	354	ARG	NE-CZ-NH2	-5.15	117.72	120.30
1	A	208	ARG	NE-CZ-NH2	-5.13	117.73	120.30
1	A	314	LYS	CD-CE-NZ	-5.12	99.92	111.70
1	A	68	ARG	NE-CZ-NH1	-5.08	117.76	120.30
1	A	186	GLU	OE1-CD-OE2	-5.04	117.25	123.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	74[A]	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3123	0	3022	16	0
2	A	2	0	0	0	0
3	A	12	0	10	0	0
4	A	512	0	0	9	3
All	All	3649	0	3032	16	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328[B]:GLU:OE1	4:A:501:HOH:O	2.12	0.68
1:A:328[A]:GLU:OE1	4:A:502:HOH:O	2.12	0.68
1:A:376:ASP:OD1	4:A:503:HOH:O	2.16	0.61
1:A:258:LEU:HD21	4:A:541:HOH:O	2.02	0.59
1:A:41[B]:ARG:NH1	1:A:45:GLU:OE2	2.42	0.52
1:A:374:ARG:NE	4:A:511:HOH:O	2.43	0.51
1:A:368[B]:ARG:NH1	4:A:515:HOH:O	2.48	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:LYS:HE2	4:A:660:HOH:O	2.16	0.45
1:A:32:ARG:HG2	1:A:296[A]:PHE:HE2	1.82	0.44
1:A:205:ARG:HD2	4:A:854:HOH:O	2.21	0.41
1:A:74[A]:ARG:HH11	1:A:74[A]:ARG:HD2	1.85	0.41
1:A:374:ARG:CZ	4:A:511:HOH:O	2.68	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:503:HOH:O	4:A:719:HOH:O[4_554]	0.96	1.24
4:A:510:HOH:O	4:A:853:HOH:O[2_555]	1.70	0.50
4:A:947:HOH:O	4:A:961:HOH:O[3_554]	2.14	0.06

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	395/388 (102%)	383 (97%)	11 (3%)	1 (0%)	46 41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	313/304 (103%)	312 (100%)	1 (0%)	94 96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	362	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GLC	A	403	2	12,12,12	0.98	1 (8%)	17,17,17	1.77	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	403	2	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	GLC	O2-C2	2.61	1.49	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	403	GLC	O3-C3-C4	-2.81	104.02	110.36
3	A	403	GLC	O2-C2-C3	2.39	115.75	110.36
3	A	403	GLC	O4-C4-C5	2.98	117.09	109.23
3	A	403	GLC	O3-C3-C2	4.14	119.69	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	387/388 (99%)	-0.69	2 (0%) 91 92	3, 6, 19, 53	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	388	GLY	4.7
1	A	2	ASN	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
3	GLC	A	403	12/12	0.98	0.08	1.37	5,6,7,7	0
2	MN	A	402	1/1	1.00	0.08	0.64	6,6,6,6	0
2	MN	A	401	1/1	0.99	0.06	-	6,6,6,6	0

6.5 Other polymers (i)

There are no such residues in this entry.