



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 9, 2017 – 03:24 PM EST

PDB ID : 1ZDH  
Title : MS2 COAT PROTEIN/RNA COMPLEX  
Authors : Valegard, K.; Van Den Worm, S.; Liljas, L.  
Deposited on : 1996-09-24  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)
RNA backbone	2183	1069 (3.10-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	19	<div><div>63%</div><div><div>16%</div><div>53%</div><div>32%</div></div></div>
1	S	19	<div><div>68%</div><div><div>47%</div><div>21%</div><div>32%</div></div></div>
2	A	129	<div><div>3%</div><div><div>80%</div><div>16%</div><div>.</div></div></div>
2	B	129	<div><div>2%</div><div><div>87%</div><div>9%</div><div>5%</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	129	<div><div></div><div>4%</div><div>84%</div><div>13%</div><div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3613 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(\*AP\*CP\*AP\*UP\*GP\*AP\*GP\*GP\*AP\*UP\*CP\*AP\*CP\*CP\*CP\*AP\*U P\*GP\*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	R	13	Total	C	N	O	P	17	0	0
			277	124	51	89	13			
1	S	13	Total	C	N	O	P	17	0	0
			274	124	51	87	12			

- Molecule 2 is a protein called PROTEIN (BACTERIOPHAGE MS2 COAT PROTEIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	129	Total	C	N	O	S	0	0	0
			965	606	165	190	4			
2	B	129	Total	C	N	O	S	0	0	0
			965	606	165	190	4			
2	C	129	Total	C	N	O	S	0	0	0
			965	606	165	190	4			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	46	Total	O	0	0
			46	46		
3	B	68	Total	O	0	0
			68	68		
3	C	49	Total	O	0	0
			49	49		
3	R	3	Total	O	0	0
			3	3		
3	S	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (5'-R(\*AP\*CP\*AP\*UP\*GP\*AP\*GP\*GP\*AP\*UP\*CP\*AP\*CP\*CP\*CP\*AP\*U P\*GP\*U)-3')




- Molecule 1: RNA (5'-R(\*AP\*CP\*AP\*UP\*GP\*AP\*GP\*GP\*AP\*UP\*CP\*AP\*CP\*CP\*CP\*AP\*U P\*GP\*U)-3')




- Molecule 2: PROTEIN (BACTERIOPHAGE MS2 COAT PROTEIN)



- Molecule 2: PROTEIN (BACTERIOPHAGE MS2 COAT PROTEIN)



- Molecule 2: PROTEIN (BACTERIOPHAGE MS2 COAT PROTEIN)



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	288.00Å 288.00Å 653.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.70 10.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	82.0 (10.00-2.70) 77.2 (10.00-2.70)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.28 (at 2.71Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.186 , (Not available) 0.164 , 0.170	Depositor DCC
$R_{free}$ test set	10642 reflections (5.23%)	DCC
Wilson B-factor (Å <sup>2</sup> )	29.3	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.24 , 72.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.18	EDS
Total number of atoms	3613	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	R	0.94	0/309	0.77	0/479
1	S	1.05	1/306 (0.3%)	0.95	1/475 (0.2%)
2	A	0.82	0/982	0.96	4/1337 (0.3%)
2	B	0.80	0/982	0.89	2/1337 (0.1%)
2	C	0.73	0/982	0.89	2/1337 (0.1%)
All	All	0.83	1/3561 (0.0%)	0.90	9/4965 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	R	0	1
1	S	0	3
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	S	4	U	N1-C2	6.92	1.44	1.38

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	SER	N-CA-C	-8.40	88.33	111.00
1	S	16	A	C2'-C3'-O3'	6.92	124.77	113.70
2	B	2	SER	N-CA-C	-6.43	93.64	111.00
2	C	13	GLY	N-CA-C	-5.85	98.48	113.10
2	A	38	ARG	NE-CZ-NH2	-5.81	117.40	120.30
2	A	48	VAL	CB-CA-C	-5.61	100.73	111.40
2	C	48	VAL	CB-CA-C	-5.35	101.23	111.40
2	B	48	VAL	CB-CA-C	-5.30	101.33	111.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	38	ARG	NE-CZ-NH1	5.03	122.81	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	R	6	A	Sidechain
1	S	14	C	Sidechain
1	S	5	G	Sidechain
1	S	7	G	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	277	0	142	7	0
1	S	274	0	143	26	0
2	A	965	0	964	11	0
2	B	965	0	964	10	0
2	C	965	0	964	10	0
3	A	46	0	0	1	0
3	B	68	0	0	3	0
3	C	49	0	0	1	0
3	R	3	0	0	0	0
3	S	1	0	0	0	0
All	All	3613	0	3177	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:20:VAL:HG13	2:C:32:TRP:HB3	1.45	0.98
2:A:20:VAL:HG13	2:A:32:TRP:HB3	1.48	0.94

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:20:VAL:HG13	2:B:32:TRP:HB3	1.48	0.94
1:S:15:C:H2'	1:S:16:A:C8	2.17	0.78
1:S:13:C:C2	1:S:14:C:C6	2.76	0.73
1:S:14:C:O2'	1:S:15:C:H5'	1.92	0.70
1:S:8:G:O2'	1:S:9:A:H5'	1.94	0.67
1:S:12:A:N7	2:C:61:LYS:HD2	2.10	0.66
1:S:15:C:H2'	1:S:16:A:H8	1.59	0.66
2:C:38:ARG:HD3	3:C:149:HOH:O	1.95	0.66
1:S:13:C:C4	1:S:14:C:C4	2.84	0.65
1:S:13:C:N3	1:S:14:C:C4	2.64	0.65
1:S:13:C:H2'	1:S:14:C:C6	2.32	0.65
1:S:13:C:H2'	1:S:14:C:H6	1.62	0.62
2:A:49:ARG:NH2	2:A:57:LYS:HD3	2.15	0.61
1:S:13:C:N3	1:S:14:C:C5	2.69	0.61
2:B:49:ARG:NH1	3:B:179:HOH:O	2.34	0.61
1:R:7:G:N2	1:R:15:C:H1'	2.16	0.61
2:A:97:THR:HG22	2:A:99:SER:H	1.66	0.59
1:R:9:A:H2'	1:R:11:C:C5	2.36	0.59
1:S:9:A:H2'	1:S:11:C:C5	2.37	0.59
1:S:13:C:C4	1:S:14:C:C5	2.90	0.59
1:R:9:A:O2'	1:R:10:U:H2'	2.04	0.58
1:S:5:G:C2	1:S:16:A:C2	2.94	0.56
2:B:49:ARG:NH2	2:B:57:LYS:HE3	2.22	0.55
1:S:13:C:N4	1:S:14:C:N4	2.55	0.54
1:S:13:C:C2	1:S:14:C:C5	2.97	0.53
2:B:83:ARG:HD2	3:B:152:HOH:O	2.10	0.52
2:A:8:VAL:HA	2:A:19:THR:HA	1.91	0.52
2:A:31:GLU:HG2	2:A:32:TRP:N	2.25	0.51
2:A:97:THR:HG22	2:A:99:SER:N	2.26	0.50
1:S:7:G:OP2	1:S:7:G:H4'	2.13	0.49
1:S:8:G:C2	1:S:9:A:C4	3.01	0.49
2:B:113:LYS:HG2	2:B:114:ASP:N	2.28	0.48
1:S:8:G:C4	1:S:9:A:C8	3.01	0.48
2:A:60:ILE:HB	2:A:88:MET:HG2	1.94	0.48
1:R:14:C:H2'	1:R:15:C:H6	1.79	0.48
1:R:5:G:C2	1:R:16:A:C2	3.02	0.47
1:R:14:C:O2'	1:R:15:C:H5'	2.16	0.47
2:A:56:ARG:CD	2:B:38:ARG:HD3	2.46	0.46
2:A:98:ASN:O	2:A:102:GLU:HG3	2.16	0.46
1:S:9:A:O2'	1:S:10:U:H2'	2.16	0.46
2:B:60:ILE:HD12	2:B:88:MET:HE2	1.99	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:8:G:O2'	1:R:9:A:H5'	2.17	0.45
1:S:6:A:H5''	1:S:7:G:C5'	2.47	0.44
2:B:3:ASN:H	2:B:3:ASN:ND2	2.16	0.43
2:C:40:GLN:HB3	2:C:67:VAL:HG22	1.99	0.42
2:C:30:ALA:O	2:C:45:THR:HA	2.18	0.42
2:C:77:LEU:HA	2:C:78:PRO:HD3	1.88	0.42
1:S:11:C:H2'	2:C:85:TYR:CE2	2.55	0.42
1:S:12:A:C8	2:C:61:LYS:HD2	2.54	0.42
2:B:56:ARG:NH2	3:B:161:HOH:O	2.49	0.42
2:C:70:GLN:O	2:C:76:GLU:HA	2.20	0.42
1:S:5:G:C6	1:S:7:G:N7	2.87	0.42
1:S:8:G:H2'	1:S:9:A:H8	1.85	0.41
2:A:75:VAL:HA	3:A:171:HOH:O	2.19	0.41
1:S:8:G:N1	1:S:9:A:C5	2.89	0.41
2:B:31:GLU:CD	2:B:38:ARG:HH22	2.23	0.41
2:A:50:GLN:HG3	2:A:52:SER:O	2.21	0.41
2:C:60:ILE:HB	2:C:88:MET:HG3	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	127/129 (98%)	119 (94%)	6 (5%)	2 (2%)	12	30
2	B	127/129 (98%)	123 (97%)	3 (2%)	1 (1%)	24	51
2	C	127/129 (98%)	121 (95%)	6 (5%)	0	100	100
All	All	381/387 (98%)	363 (95%)	15 (4%)	3 (1%)	24	51

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	2	SER
2	B	2	SER
2	A	14	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	106/106 (100%)	97 (92%)	9 (8%)	13	30
2	B	106/106 (100%)	99 (93%)	7 (7%)	21	45
2	C	106/106 (100%)	100 (94%)	6 (6%)	25	53
All	All	318/318 (100%)	296 (93%)	22 (7%)	19	43

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	4	PHE
2	A	20	VAL
2	A	25	PHE
2	A	33	ILE
2	A	35	SER
2	A	49	ARG
2	A	52	SER
2	A	79	VAL
2	A	97	THR
2	B	3	ASN
2	B	15	THR
2	B	20	VAL
2	B	48	VAL
2	B	57	LYS
2	B	77	LEU
2	B	114	ASP
2	C	20	VAL
2	C	25	PHE
2	C	35	SER
2	C	38	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	C	48	VAL
2	C	54	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	55	ASN
2	B	3	ASN
2	B	36	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	12/19 (63%)	0	0
1	S	12/19 (63%)	0	0
All	All	24/38 (63%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	R	13/19 (68%)	5.91	12 (92%) <b>0</b> <b>0</b>	36, 52, 68, 84	13 (100%)
1	S	13/19 (68%)	5.37	13 (100%) <b>0</b> <b>0</b>	44, 48, 68, 69	13 (100%)
2	A	129/129 (100%)	0.23	4 (3%) 52 52	4, 12, 40, 64	0
2	B	129/129 (100%)	0.04	3 (2%) 64 64	3, 10, 37, 59	0
2	C	129/129 (100%)	0.36	5 (3%) 43 43	6, 14, 37, 76	0
All	All	413/425 (97%)	0.55	37 (8%) <b>12</b> <b>9</b>	3, 12, 54, 84	26 (6%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	S	4	U	9.0
1	R	5	G	8.5
1	R	15	C	8.5
1	R	6	A	8.4
1	S	15	C	8.3
1	R	4	U	7.6
1	R	14	C	7.6
1	S	16	A	7.5
1	S	5	G	7.5
1	R	7	G	6.8
1	R	13	C	6.6
2	C	14	GLY	6.5
1	R	16	A	6.4
1	R	8	G	6.4
1	S	14	C	6.3
1	S	6	A	5.2
2	C	1	ALA	4.9
1	S	10	U	4.5
1	S	13	C	4.5
2	A	15	THR	4.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	S	7	G	4.0
2	B	1	ALA	3.8
1	S	11	C	3.7
2	C	15	THR	3.6
2	A	14	GLY	3.4
1	S	9	A	3.3
1	R	11	C	3.2
1	R	9	A	3.1
1	S	12	A	3.1
1	S	8	G	3.0
2	B	15	THR	2.9
2	A	53	ALA	2.7
1	R	10	U	2.3
2	C	13	GLY	2.1
2	A	70	GLN	2.1
2	C	115	GLY	2.1
2	B	53	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.