



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:28 PM GMT

PDB ID : 3ZIX  
Title : Clostridium perfringens Enterotoxin with the N-terminal 37 residues deleted  
Authors : Yelland, T.; Naylor, C.E.; Savva, C.G.; Basak, A.K.  
Deposited on : 2013-01-14  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	<div> <div>2%</div> <div>95%</div> <div>5%</div> </div>
1	B	286	<div> <div>2%</div> <div>95%</div> <div>5%</div> </div>
1	C	286	<div> <div>7%</div> <div>92%</div> <div>7%</div> </div>
1	D	286	<div> <div>6%</div> <div>96%</div> <div>• •</div> </div>
1	E	286	<div> <div>8%</div> <div>94%</div> <div>6%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	286	 % 93% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	P6G	A	400	-	-	-	X
2	P6G	A	401	-	-	-	X
2	P6G	A	402	-	-	-	X
2	P6G	B	400	-	-	-	X
2	P6G	C	400	-	-	-	X
2	P6G	C	401	-	-	-	X
2	P6G	F	400	-	-	-	X
2	P6G	F	401	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEAT-LABILE ENTEROTOXIN B CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	6	0
			2251	1427	363	458	3			
1	B	286	Total	C	N	O	S	0	4	0
			2228	1415	357	452	4			
1	C	286	Total	C	N	O	S	0	4	0
			2237	1419	363	452	3			
1	D	286	Total	C	N	O	S	0	7	0
			2236	1419	358	456	3			
1	E	286	Total	C	N	O	S	0	5	0
			2220	1412	357	448	3			
1	F	286	Total	C	N	O	S	0	4	0
			2243	1420	365	455	3			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	34	GLY	-	EXPRESSION TAG	UNP P01558
A	35	ALA	-	EXPRESSION TAG	UNP P01558
A	36	MET	-	EXPRESSION TAG	UNP P01558
A	37	GLY	-	EXPRESSION TAG	UNP P01558
B	34	GLY	-	EXPRESSION TAG	UNP P01558
B	35	ALA	-	EXPRESSION TAG	UNP P01558
B	36	MET	-	EXPRESSION TAG	UNP P01558
B	37	GLY	-	EXPRESSION TAG	UNP P01558
C	34	GLY	-	EXPRESSION TAG	UNP P01558
C	35	ALA	-	EXPRESSION TAG	UNP P01558
C	36	MET	-	EXPRESSION TAG	UNP P01558
C	37	GLY	-	EXPRESSION TAG	UNP P01558
D	34	GLY	-	EXPRESSION TAG	UNP P01558
D	35	ALA	-	EXPRESSION TAG	UNP P01558
D	36	MET	-	EXPRESSION TAG	UNP P01558
D	37	GLY	-	EXPRESSION TAG	UNP P01558
E	34	GLY	-	EXPRESSION TAG	UNP P01558

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	35	ALA	-	EXPRESSION TAG	UNP P01558
E	36	MET	-	EXPRESSION TAG	UNP P01558
E	37	GLY	-	EXPRESSION TAG	UNP P01558
F	34	GLY	-	EXPRESSION TAG	UNP P01558
F	35	ALA	-	EXPRESSION TAG	UNP P01558
F	36	MET	-	EXPRESSION TAG	UNP P01558
F	37	GLY	-	EXPRESSION TAG	UNP P01558

- Molecule 2 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: C<sub>12</sub>H<sub>26</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			19	12	7		
2	A	1	Total	C	O	0	0
			14	9	5		
2	A	1	Total	C	O	0	0
			10	6	4		
2	B	1	Total	C	O	0	0
			17	11	6		
2	B	1	Total	C	O	0	0
			11	7	4		
2	C	1	Total	C	O	0	0
			16	10	6		
2	C	1	Total	C	O	0	0
			8	5	3		
2	D	1	Total	C	O	0	0
			16	10	6		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			10	6	4		
2	E	1	Total	C	O	0	0
			14	9	5		
2	E	1	Total	C	O	0	0
			13	8	5		
2	E	1	Total	C	O	0	0
			10	6	4		
2	F	1	Total	C	O	0	0
			19	12	7		
2	F	1	Total	C	O	0	0
			8	5	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	305	Total	O	0	0
			305	305		
3	B	314	Total	O	0	0
			314	314		
3	C	266	Total	O	0	0
			266	266		
3	D	272	Total	O	0	0
			272	272		
3	E	268	Total	O	0	0
			268	268		
3	F	336	Total	O	0	0
			336	336		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HEAT-LABILE ENTEROTOXIN B CHAIN




- Molecule 1: HEAT-LABILE ENTEROTOXIN B CHAIN



- Molecule 1: HEAT-LABILE ENTEROTOXIN B CHAIN



- Molecule 1: HEAT-LABILE ENTEROTOXIN B CHAIN




- Molecule 1: HEAT-LABILE ENTEROTOXIN B CHAIN





● Molecule 1: HEAT-LABILE ENTEROTOXIN B CHAIN





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	191.68Å 128.29Å 137.20Å 90.00° 133.81° 90.00°	Depositor
Resolution (Å)	47.03 – 1.90 47.03 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.8 (47.03-1.90) 98.8 (47.03-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.08 (at 1.90Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, $R_{free}$	0.176 , 0.197 0.180 , 0.200	Depositor DCC
$R_{free}$ test set	9299 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.8	Xtriage
Anisotropy	0.105	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 62.2	EDS
Estimated twinning fraction	0.005 for h+2*l,k,-h-l 0.012 for h,-k,-h-l 0.038 for -h-2*l,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 185657 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	15361	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: P6G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/2315	0.66	0/3141
1	B	0.54	0/2284	0.64	0/3105
1	C	0.51	0/2292	0.65	0/3112
1	D	0.51	0/2301	0.66	0/3129
1	E	0.52	0/2278	0.66	0/3099
1	F	0.55	0/2299	0.64	0/3121
All	All	0.52	0/13769	0.65	0/18707

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2251	0	2191	8	0
1	B	2228	0	2158	10	0
1	C	2237	0	2181	15	0
1	D	2236	0	2168	10	0
1	E	2220	0	2160	11	0
1	F	2243	0	2189	17	0
2	A	43	0	55	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	28	0	34	4	0
2	C	24	0	30	0	0
2	D	26	0	34	0	0
2	E	37	0	46	1	0
2	F	27	0	35	2	0
3	A	305	0	0	2	0
3	B	314	0	0	1	0
3	C	266	0	0	0	0
3	D	272	0	0	0	0
3	E	268	0	0	2	0
3	F	336	0	0	0	0
All	All	15361	0	13281	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:110[A]:GLU:HG3	1:F:110[A]:GLU:HG3	1.46	0.96
1:A:110[A]:GLU:HG2	1:C:110[A]:GLU:HG3	1.47	0.92
1:F:62:ILE:HG21	1:F:171:LEU:HD21	1.63	0.81
1:E:263[B]:ILE:HD11	1:E:287:VAL:HG21	1.63	0.80
1:B:263:ILE:HD11	1:B:287:VAL:HG21	1.73	0.70
1:C:110[B]:GLU:CD	1:C:110[B]:GLU:H	1.96	0.67
1:F:78:SER:HA	1:F:111[B]:GLN:HG3	1.81	0.63
1:E:49:GLY:HA3	1:E:188:LEU:HG	1.80	0.62
1:E:264:TYR:HE1	1:E:299[B]:VAL:CG1	2.13	0.62
1:C:263:ILE:HD11	1:C:287:VAL:HG21	1.80	0.62
1:D:263:ILE:HD11	1:D:287:VAL:HG21	1.83	0.60
1:D:110[A]:GLU:CG	1:F:110[A]:GLU:HG3	2.27	0.60
1:C:120:THR:HG21	1:C:165:LYS:HD3	1.85	0.58
1:A:49:GLY:HA3	1:A:188:LEU:HG	1.86	0.57
1:E:110:GLU:HG2	3:E:2125:HOH:O	2.05	0.57
1:E:264:TYR:HE1	1:E:299[B]:VAL:HG11	1.71	0.56
1:E:110:GLU:CG	3:E:2125:HOH:O	2.54	0.56
1:B:49:GLY:HA3	1:B:188:LEU:HG	1.88	0.54
1:D:49:GLY:HA3	1:D:188:LEU:HG	1.90	0.54
1:B:82:SER:HA	2:B:400:P6G:H172	1.89	0.53
1:A:78:SER:HA	1:A:111[A]:GLN:HG3	1.92	0.52
1:B:263:ILE:CD1	1:B:287:VAL:HG21	2.38	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:120:THR:HG21	1:E:165:LYS:HD3	1.92	0.51
1:B:81:VAL:O	2:B:400:P6G:H172	2.11	0.50
1:F:280:ASP:HA	2:F:401:P6G:H62	1.93	0.50
1:F:79:LYS:NZ	1:F:110[A]:GLU:OE2	2.39	0.49
1:F:49:GLY:HA3	1:F:188:LEU:HG	1.93	0.49
1:D:78:SER:HA	1:D:111[A]:GLN:HG3	1.93	0.49
1:B:82:SER:HB2	2:B:400:P6G:H122	1.94	0.49
1:C:122:ALA:HB2	1:C:128:VAL:HG22	1.94	0.49
1:C:79:LYS:HE3	1:C:110[A]:GLU:OE2	2.13	0.48
1:B:78:SER:HA	1:B:111:GLN:HG3	1.94	0.48
1:D:166[A]:THR:HG21	1:D:178:SER:OG	2.13	0.47
1:E:81:VAL:O	2:E:400:P6G:H152	2.14	0.47
1:C:263:ILE:CD1	1:C:287:VAL:HG21	2.45	0.47
1:C:287:VAL:HG23	1:C:289:ILE:HD12	1.97	0.47
1:D:110[A]:GLU:HG3	1:F:110[A]:GLU:CG	2.32	0.46
1:F:215:LEU:O	1:F:221:GLY:HA2	2.15	0.46
1:A:192:SER:HB2	3:A:2018:HOH:O	2.16	0.46
1:D:215:LEU:O	1:D:221:GLY:HA2	2.17	0.45
1:A:110[B]:GLU:HG2	3:A:2139:HOH:O	2.17	0.45
1:D:263:ILE:CD1	1:D:287:VAL:HG21	2.46	0.45
1:C:78:SER:HA	1:C:111[A]:GLN:HG3	1.99	0.45
1:E:78:SER:HA	1:E:111:GLN:HG3	1.99	0.44
1:B:110:GLU:HG2	3:B:2144:HOH:O	2.16	0.44
1:B:215:LEU:O	1:B:221:GLY:HA2	2.17	0.44
1:A:215:LEU:O	1:A:221:GLY:HA2	2.18	0.44
1:C:128:VAL:HG12	1:C:167:SER:HB3	1.99	0.44
1:C:215:LEU:O	1:C:221:GLY:HA2	2.17	0.44
1:A:43:VAL:HG21	1:A:103:GLY:HA3	2.01	0.43
1:E:262:ASN:HB2	1:E:299[B]:VAL:CG1	2.49	0.43
1:E:264:TYR:HE1	1:E:299[B]:VAL:HG12	1.82	0.43
1:A:110[A]:GLU:HG2	1:C:110[A]:GLU:CG	2.34	0.42
1:F:263:ILE:HD11	1:F:287:VAL:HG21	2.02	0.42
1:C:43:VAL:HG21	1:C:103:GLY:HA3	2.02	0.42
1:B:52:LEU:HD22	2:B:400:P6G:H121	2.01	0.42
1:F:62:ILE:CG2	1:F:171:LEU:HD21	2.40	0.41
1:F:263:ILE:CD1	1:F:287:VAL:HG21	2.51	0.41
1:D:110[B]:GLU:OE1	1:F:110[B]:GLU:HG2	2.21	0.41
1:C:264:TYR:HB2	1:C:297[B]:VAL:HG13	2.01	0.41
1:F:81:VAL:O	2:F:400:P6G:H51	2.20	0.41
1:F:257:LYS:HG3	1:F:258:ILE:HG23	2.02	0.40
1:F:43:VAL:HG21	1:F:103:GLY:HA3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/286 (101%)	285 (99%)	4 (1%)	0	100	100
1	B	288/286 (101%)	283 (98%)	5 (2%)	0	100	100
1	C	288/286 (101%)	285 (99%)	3 (1%)	0	100	100
1	D	291/286 (102%)	288 (99%)	3 (1%)	0	100	100
1	E	289/286 (101%)	285 (99%)	4 (1%)	0	100	100
1	F	288/286 (101%)	284 (99%)	4 (1%)	0	100	100
All	All	1733/1716 (101%)	1710 (99%)	23 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/247 (102%)	246 (98%)	5 (2%)	63	57
1	B	246/247 (100%)	243 (99%)	3 (1%)	78	76
1	C	249/247 (101%)	241 (97%)	8 (3%)	46	35
1	D	249/247 (101%)	244 (98%)	5 (2%)	63	57
1	E	247/247 (100%)	242 (98%)	5 (2%)	63	57
1	F	251/247 (102%)	246 (98%)	5 (2%)	63	57

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	1493/1482 (101%)	1462 (98%)	31 (2%)	66	55

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60[A]	SER
1	A	60[B]	SER
1	A	171	LEU
1	A	292	ASP
1	A	307	SER
1	B	256	SER
1	B	269	ASN
1	B	292	ASP
1	C	111[A]	GLN
1	C	111[B]	GLN
1	C	188	LEU
1	C	256	SER
1	C	258	ILE
1	C	289	ILE
1	C	292	ASP
1	C	307	SER
1	D	110[A]	GLU
1	D	110[B]	GLU
1	D	111[A]	GLN
1	D	111[B]	GLN
1	D	292	ASP
1	E	249	GLN
1	E	269	ASN
1	E	275	GLU
1	E	292	ASP
1	E	307	SER
1	F	111[A]	GLN
1	F	111[B]	GLN
1	F	171	LEU
1	F	292	ASP
1	F	307	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	269	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	P6G	A	400	-	18,18,18	0.50	0	17,17,17	0.46	0
2	P6G	A	401	-	13,13,18	0.48	0	12,12,17	0.25	0
2	P6G	A	402	-	9,9,18	0.48	0	8,8,17	0.28	0
2	P6G	B	400	-	16,16,18	0.53	0	15,15,17	0.64	0
2	P6G	B	401	-	10,10,18	0.46	0	9,9,17	0.27	0
2	P6G	C	400	-	15,15,18	0.52	0	14,14,17	0.44	0
2	P6G	C	401	-	7,7,18	0.49	0	6,6,17	0.18	0
2	P6G	D	400	-	15,15,18	0.49	0	14,14,17	0.55	0
2	P6G	D	401	-	9,9,18	0.48	0	8,8,17	0.31	0
2	P6G	E	400	-	13,13,18	0.55	0	12,12,17	0.23	0
2	P6G	E	401	-	12,12,18	0.50	0	11,11,17	0.24	0
2	P6G	E	403	-	9,9,18	0.45	0	8,8,17	0.31	0
2	P6G	F	400	-	18,18,18	0.43	0	17,17,17	0.54	0
2	P6G	F	401	-	7,7,18	0.49	0	6,6,17	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P6G	A	400	-	-	0/16/16/16	0/0/0/0
2	P6G	A	401	-	-	0/11/11/16	0/0/0/0
2	P6G	A	402	-	-	0/7/7/16	0/0/0/0
2	P6G	B	400	-	-	0/14/14/16	0/0/0/0
2	P6G	B	401	-	-	0/8/8/16	0/0/0/0
2	P6G	C	400	-	-	0/13/13/16	0/0/0/0
2	P6G	C	401	-	-	0/5/5/16	0/0/0/0
2	P6G	D	400	-	-	0/13/13/16	0/0/0/0
2	P6G	D	401	-	-	0/7/7/16	0/0/0/0
2	P6G	E	400	-	-	0/11/11/16	0/0/0/0
2	P6G	E	401	-	-	0/10/10/16	0/0/0/0
2	P6G	E	403	-	-	0/7/7/16	0/0/0/0
2	P6G	F	400	-	-	0/16/16/16	0/0/0/0
2	P6G	F	401	-	-	0/5/5/16	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	400	P6G	4	0
2	E	400	P6G	1	0
2	F	400	P6G	1	0
2	F	401	P6G	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	286/286 (100%)	-0.06	7 (2%) 62 66	25, 45, 68, 81	0
1	B	286/286 (100%)	-0.07	6 (2%) 67 70	23, 40, 66, 90	0
1	C	286/286 (100%)	0.25	21 (7%) 18 20	24, 43, 79, 101	0
1	D	286/286 (100%)	0.07	18 (6%) 23 26	26, 45, 76, 96	0
1	E	286/286 (100%)	0.16	22 (7%) 16 18	23, 44, 74, 103	0
1	F	286/286 (100%)	-0.07	3 (1%) 84 86	22, 36, 58, 83	0
All	All	1716/1716 (100%)	0.05	77 (4%) 37 40	22, 42, 72, 103	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	LEU	5.7
1	C	66	ASN	4.9
1	E	249	GLN	4.8
1	C	319	PHE	4.8
1	C	219	PRO	4.6
1	B	219	PRO	4.5
1	E	127	TYR	4.3
1	C	62	ILE	4.3
1	C	127	TYR	4.2
1	C	65	PRO	4.2
1	C	124	PRO	4.0
1	C	68	THR	3.8
1	D	168	ALA	3.8
1	C	69	GLY	3.7
1	B	319	PHE	3.7
1	D	219	PRO	3.6
1	C	122	ALA	3.6
1	C	120	THR	3.6
1	C	123	GLY	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	221	GLY	3.5
1	D	169	ASP	3.5
1	D	122	ALA	3.4
1	D	128	VAL	3.4
1	C	168	ALA	3.4
1	D	310	TYR	3.3
1	C	125	ASN	3.3
1	E	69	GLY	3.2
1	E	247	THR	3.2
1	E	319	PHE	3.2
1	D	173	ASN	3.2
1	E	68	THR	3.1
1	D	124	PRO	3.1
1	F	219	PRO	3.1
1	E	220	ALA	3.1
1	E	307	SER	3.1
1	E	306	TYR	3.0
1	E	66	ASN	3.0
1	B	247	THR	3.0
1	E	248	GLY	2.9
1	D	64	ASN	2.8
1	F	310	TYR	2.8
1	D	170	SER	2.8
1	B	307	SER	2.8
1	C	70	THR	2.7
1	D	306	TYR	2.6
1	E	123	GLY	2.6
1	B	220	ALA	2.6
1	E	218	ASN	2.5
1	C	119	SER	2.5
1	C	308	GLY	2.5
1	D	127	TYR	2.5
1	E	125	ASN	2.4
1	E	36	MET	2.4
1	A	220	ALA	2.4
1	E	293	ALA	2.4
1	A	310	TYR	2.4
1	A	125	ASN	2.3
1	E	37	GLY	2.3
1	E	124	PRO	2.3
1	C	67	GLU	2.3
1	D	62	ILE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	169	ASP	2.3
1	B	221	GLY	2.3
1	E	171	LEU	2.3
1	D	220	ALA	2.3
1	D	224	TYR	2.2
1	E	219	PRO	2.2
1	C	171	LEU	2.2
1	F	221	GLY	2.2
1	C	221	GLY	2.1
1	A	307	SER	2.1
1	A	169	ASP	2.1
1	D	36	MET	2.1
1	D	248	GLY	2.1
1	E	65	PRO	2.1
1	D	221	GLY	2.1
1	A	219	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	P6G	F	400	19/19	0.85	0.52	20.91	27,36,43,48	19
2	P6G	F	401	8/19	0.75	0.17	9.66	61,64,72,73	0
2	P6G	C	400	16/19	0.90	0.16	4.91	57,61,67,67	0
2	P6G	B	400	17/19	0.84	0.16	3.44	45,52,61,61	0
2	P6G	A	400	19/19	0.91	0.14	2.33	45,59,68,71	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	P6G	A	401	14/19	0.77	0.18	2.12	74,77,80,81	0
2	P6G	A	402	10/19	0.88	0.17	2.00	69,74,84,85	0
2	P6G	C	401	8/19	0.84	0.14	2.00	57,59,62,63	0
2	P6G	E	403	10/19	0.88	0.18	1.97	53,59,62,63	0
2	P6G	E	400	14/19	0.92	0.12	0.24	49,53,63,65	0
2	P6G	E	401	13/19	0.86	0.11	0.18	57,62,65,66	0
2	P6G	B	401	11/19	0.84	0.10	0.07	60,62,65,65	0
2	P6G	D	401	10/19	0.78	0.12	0.03	74,75,76,77	0
2	P6G	D	400	16/19	0.92	0.09	-0.09	45,53,70,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.