



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:05 AM GMT

PDB ID : 1ZKK
Title : Crystal structure of hSET8 in ternary complex with H4 peptide (16-24) and AdoHcy
Authors : Couture, J.-F.; Collazo, E.; Brunzelle, J.S.; Trievel, R.C.
Deposited on : 2005-05-03
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1278 (1.48-1.44)
Clashscore	102246	1336 (1.48-1.44)
Ramachandran outliers	100387	1320 (1.48-1.44)
Sidechain outliers	100360	1320 (1.48-1.44)
RSRZ outliers	91569	1279 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	<div> <div>6%</div> <div>89%</div> <div>6%</div> <div>• •</div> </div>
1	B	167	<div> <div>4%</div> <div>93%</div> <div>•</div> <div>•</div> </div>
1	C	167	<div> <div>2%</div> <div>91%</div> <div>5%</div> <div>•</div> </div>
1	D	167	<div> <div>%</div> <div>90%</div> <div>6%</div> <div>•</div> </div>
2	E	10	<div> <div>10%</div> <div>80%</div> <div>20%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	10	
2	G	10	
2	H	10	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase, H4 lysine-20 specific.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	160	Total	C	N	O	S	1	2	0
			1280	799	225	249	7			
1	B	161	Total	C	N	O	S	0	1	0
			1284	802	227	249	6			
1	C	160	Total	C	N	O	S	0	0	0
			1278	799	225	248	6			
1	D	161	Total	C	N	O	S	0	2	0
			1290	805	228	251	6			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	186	GLY	-	CLONING ARTIFACT	UNP Q9NQR1
A	187	ALA	-	CLONING ARTIFACT	UNP Q9NQR1
A	188	MET	-	CLONING ARTIFACT	UNP Q9NQR1
A	189	GLY	-	CLONING ARTIFACT	UNP Q9NQR1
B	186	GLY	-	CLONING ARTIFACT	UNP Q9NQR1
B	187	ALA	-	CLONING ARTIFACT	UNP Q9NQR1
B	188	MET	-	CLONING ARTIFACT	UNP Q9NQR1
B	189	GLY	-	CLONING ARTIFACT	UNP Q9NQR1
C	186	GLY	-	CLONING ARTIFACT	UNP Q9NQR1
C	187	ALA	-	CLONING ARTIFACT	UNP Q9NQR1
C	188	MET	-	CLONING ARTIFACT	UNP Q9NQR1
C	189	GLY	-	CLONING ARTIFACT	UNP Q9NQR1
D	186	GLY	-	CLONING ARTIFACT	UNP Q9NQR1
D	187	ALA	-	CLONING ARTIFACT	UNP Q9NQR1
D	188	MET	-	CLONING ARTIFACT	UNP Q9NQR1
D	189	GLY	-	CLONING ARTIFACT	UNP Q9NQR1

- Molecule 2 is a protein called Peptide corresponding to residues 15-24 of histone H4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	8	Total	C	N	O	0	0	0
			72	44	20	8			
2	F	9	Total	C	N	O	0	0	0
			79	47	21	11			
2	G	9	Total	C	N	O	0	0	0
			80	48	21	11			
2	H	8	Total	C	N	O	0	0	0
			71	43	20	8			

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	C	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	D	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	218	Total	O	0	0
			218	218		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	226	Total 226	O 226	0	0
4	C	265	Total 265	O 265	0	0
4	D	253	Total 253	O 253	0	0
4	E	7	Total 7	O 7	0	0
4	F	20	Total 20	O 20	0	0
4	G	8	Total 8	O 8	0	0
4	H	10	Total 10	O 10	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-lysine N-methyltransferase, H4 lysine-20 specific

Chain A: 




- Molecule 1: Histone-lysine N-methyltransferase, H4 lysine-20 specific

Chain B: 




- Molecule 1: Histone-lysine N-methyltransferase, H4 lysine-20 specific

Chain C: 




- Molecule 1: Histone-lysine N-methyltransferase, H4 lysine-20 specific

Chain D: 





- Molecule 2: Peptide corresponding to residues 15-24 of histone H4

Chain E: 



- Molecule 2: Peptide corresponding to residues 15-24 of histone H4

Chain F:  80% 10% 10%





- Molecule 2: Peptide corresponding to residues 15-24 of histone H4

Chain G:  10% 50% 30% 10% 10%



- Molecule 2: Peptide corresponding to residues 15-24 of histone H4

Chain H:  80% 20%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.96Å 45.77Å 94.44Å 89.22° 87.07° 90.72°	Depositor
Resolution (Å)	19.63 – 1.45 19.63 – 1.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.63-1.45) 95.1 (19.63-1.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.169 , 0.199 0.175 , 0.204	Depositor DCC
R_{free} test set	6198 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.1	EDS
Estimated twinning fraction	0.016 for -h,k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 123958 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6545	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.07 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3924e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1312	0.61	0/1761
1	B	0.40	0/1314	0.62	0/1763
1	C	0.42	0/1300	0.63	0/1745
1	D	0.44	0/1322	0.63	0/1775
2	E	0.45	0/72	0.73	0/93
2	F	0.52	0/79	0.74	0/102
2	G	0.44	0/80	0.81	0/104
2	H	0.49	0/71	0.80	0/91
All	All	0.43	0/5550	0.63	0/7434

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1280	0	1261	6	0
1	B	1284	0	1258	3	0
1	C	1278	0	1259	11	0
1	D	1290	0	1267	5	0
2	E	72	0	80	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	79	0	83	1	0
2	G	80	0	84	4	0
2	H	71	0	79	0	0
3	A	26	0	19	0	0
3	B	26	0	19	0	0
3	C	26	0	19	0	0
3	D	26	0	19	0	0
4	A	218	0	0	2	0
4	B	226	0	0	1	0
4	C	265	0	0	3	0
4	D	253	0	0	1	0
4	E	7	0	0	0	0
4	F	20	0	0	1	0
4	G	8	0	0	0	0
4	H	10	0	0	0	0
All	All	6545	0	5447	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:HIS:C	4:A:3006:HOH:O	1.68	1.26
1:C:223:ILE:HD11	1:C:231:ILE:HD11	1.48	0.94
1:C:219:LYS:HE3	1:C:233:THR:HG22	1.69	0.74
1:C:221:ASP:CB	1:C:231:ILE:HD13	2.22	0.69
1:C:221:ASP:HB3	1:C:231:ILE:HD13	1.72	0.68
1:C:235:GLN:HG3	4:C:2903:HOH:O	2.02	0.60
2:G:19:ARG:HH12	2:G:21:VAL:HG21	1.69	0.58
1:A:214:LYS:HG2	1:A:216:GLU:HG3	1.87	0.56
1:D:257:LYS:HE2	1:D:257:LYS:HA	1.89	0.54
1:C:221:ASP:HB3	1:C:231:ILE:CD1	2.36	0.53
2:G:19:ARG:NH1	2:G:21:VAL:HG21	2.24	0.52
1:B:280:LYS:HD3	4:B:2908:HOH:O	2.11	0.51
1:C:221:ASP:HB2	1:C:231:ILE:HD13	1.92	0.51
2:F:23:ARG:HB3	4:F:564:HOH:O	2.10	0.51
1:C:235:GLN:CG	4:C:2903:HOH:O	2.57	0.49
1:D:306[A]:GLN:HG2	1:D:308:LYS:HE3	1.94	0.48
1:B:207:ASP:O	1:B:211:GLU:HG3	2.15	0.47
1:B:276:GLN:O	1:B:309:LEU:HD11	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:219:LYS:HE3	1:C:233:THR:CG2	2.42	0.46
1:D:259:GLU:CD	4:D:2912:HOH:O	2.53	0.46
1:A:261:LEU:HA	1:A:264:GLN:HE21	1.81	0.45
1:A:251:GLU:HG3	4:A:2875:HOH:O	2.17	0.44
1:D:210:ILE:HD11	1:D:319:ILE:HG21	2.00	0.43
1:C:325:ASP:HB3	4:C:2903:HOH:O	2.19	0.42
1:D:252:ILE:HG12	1:D:256:LYS:HD2	2.01	0.42
1:C:231:ILE:N	1:C:231:ILE:HD12	2.35	0.41
2:G:23:ARG:O	2:G:24:ASP:HB2	2.21	0.41
1:A:309:LEU:C	1:A:309:LEU:HD13	2.42	0.40
1:A:222:LEU:HD21	2:G:19:ARG:NH1	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/167 (96%)	157 (98%)	3 (2%)	0	100	100
1	B	160/167 (96%)	157 (98%)	3 (2%)	0	100	100
1	C	158/167 (95%)	156 (99%)	2 (1%)	0	100	100
1	D	161/167 (96%)	159 (99%)	2 (1%)	0	100	100
2	E	6/10 (60%)	6 (100%)	0	0	100	100
2	F	7/10 (70%)	7 (100%)	0	0	100	100
2	G	7/10 (70%)	7 (100%)	0	0	100	100
2	H	6/10 (60%)	6 (100%)	0	0	100	100
All	All	665/708 (94%)	655 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/140 (99%)	134 (97%)	4 (3%)	50	13
1	B	137/140 (98%)	136 (99%)	1 (1%)	88	69
1	C	136/140 (97%)	135 (99%)	1 (1%)	88	69
1	D	138/140 (99%)	136 (99%)	2 (1%)	74	42
2	E	7/9 (78%)	7 (100%)	0	100	100
2	F	8/9 (89%)	8 (100%)	0	100	100
2	G	8/9 (89%)	7 (88%)	1 (12%)	6	0
2	H	7/9 (78%)	7 (100%)	0	100	100
All	All	579/596 (97%)	570 (98%)	9 (2%)	70	35

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	LYS
1	A	251	GLU
1	A	273	TYR
1	A	276	GLN
1	B	273	TYR
1	C	273	TYR
1	D	251	GLU
1	D	273	TYR
2	G	23	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	264	GLN
1	A	276	GLN
1	A	291	ASN
1	D	199	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SAH	A	2800	-	20,28,28	0.93	1 (5%)	19,40,40	2.44	2 (10%)
3	SAH	B	2801	-	20,28,28	0.99	2 (10%)	19,40,40	2.37	2 (10%)
3	SAH	C	2802	-	20,28,28	0.87	2 (10%)	19,40,40	2.81	3 (15%)
3	SAH	D	2803	-	20,28,28	1.00	2 (10%)	19,40,40	2.10	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	A	2800	-	-	0/7/31/31	0/3/3/3
3	SAH	B	2801	-	-	0/7/31/31	0/3/3/3
3	SAH	C	2802	-	-	0/7/31/31	0/3/3/3
3	SAH	D	2803	-	-	0/7/31/31	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2802	SAH	C2-N1	2.17	1.38	1.33
3	D	2803	SAH	C2-N1	2.25	1.38	1.33
3	B	2801	SAH	C2-N1	2.46	1.38	1.33
3	C	2802	SAH	C2-N3	2.73	1.37	1.32
3	A	2800	SAH	C2-N3	2.93	1.37	1.32
3	B	2801	SAH	C2-N3	3.18	1.37	1.32
3	D	2803	SAH	C2-N3	3.19	1.37	1.32

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2802	SAH	N3-C2-N1	-11.55	120.05	128.89
3	A	2800	SAH	N3-C2-N1	-10.09	121.17	128.89
3	B	2801	SAH	N3-C2-N1	-9.49	121.63	128.89
3	D	2803	SAH	N3-C2-N1	-8.11	122.69	128.89
3	B	2801	SAH	C4-C5-N7	-2.17	107.48	109.48
3	C	2802	SAH	C4-C5-N7	-2.06	107.58	109.48
3	A	2800	SAH	C4-C5-N7	-2.03	107.61	109.48
3	C	2802	SAH	C2-N1-C6	2.16	122.63	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	160/167 (95%)	0.39	10 (6%) 23 23	9, 14, 25, 32	0
1	B	161/167 (96%)	0.23	6 (3%) 45 46	8, 13, 21, 25	0
1	C	160/167 (95%)	0.07	4 (2%) 61 61	7, 13, 22, 26	0
1	D	161/167 (96%)	0.06	2 (1%) 81 82	7, 12, 22, 28	0
2	E	8/10 (80%)	0.33	1 (12%) 5 5	11, 15, 21, 25	0
2	F	9/10 (90%)	0.27	0 100 100	10, 12, 17, 22	0
2	G	9/10 (90%)	0.85	1 (11%) 7 6	10, 16, 27, 30	0
2	H	8/10 (80%)	0.02	0 100 100	7, 13, 18, 20	0
All	All	676/708 (95%)	0.20	24 (3%) 46 47	7, 13, 22, 32	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	266	PRO	6.6
1	A	261	LEU	4.6
1	A	352	HIS	3.7
1	A	267	SER	3.7
2	G	24	ASP	3.5
1	B	192	ARG	3.0
1	B	264	GLN	3.0
1	A	264	GLN	2.9
1	A	263	ALA	2.9
1	D	196	ALA	2.8
1	A	268	THR	2.7
1	A	332	LEU	2.6
1	D	267	SER	2.6
1	C	264	GLN	2.5
1	A	265	ASP	2.5
1	A	260	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	261	LEU	2.3
2	E	16	LYS	2.2
1	C	203	ARG	2.1
1	B	194	SER	2.1
1	B	257	LYS	2.1
1	B	352	HIS	2.1
1	C	345	GLU	2.1
1	C	257	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SAH	D	2803	26/26	0.98	0.07	-0.66	6,7,8,8	0
3	SAH	C	2802	26/26	0.98	0.07	-0.67	7,9,10,10	0
3	SAH	A	2800	26/26	0.98	0.07	-0.70	8,9,11,11	0
3	SAH	B	2801	26/26	0.98	0.07	-0.82	7,8,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.