



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:04 AM GMT

PDB ID : 1ZKR  
Title : Crystal structure of the major cat allergen Fel d 1 (1+2)  
Authors : Kaiser, L.; Cirkovic Velickovic, T.; Adedoyin, J.; Thunberg, S.; Gronlund, H.; Gafvelin, G.; van Hage, M.; Achour, A.  
Deposited on : 2005-05-04  
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1953 (1.66-1.62)
Clashscore	102246	2091 (1.66-1.62)
Ramachandran outliers	100387	2052 (1.66-1.62)
Sidechain outliers	100360	2052 (1.66-1.62)
RSRZ outliers	91569	1955 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	<div> <div>3%</div> <div>82%</div> <div>10%</div> <div>• • 5%</div> </div>
1	B	153	<div> <div>%</div> <div>78%</div> <div>12%</div> <div>• • 5%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major allergen I polypeptide, fused chain 1, chain 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	0	0
			1129	714	179	224	12			
1	B	145	Total	C	N	O	S	0	0	0
			1125	712	178	223	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	INITIATING METHIONINE	UNP P30438
A	60	LEU	VAL	CONFLICT	UNP P30438
A	147	HIS	-	EXPRESSION TAG	UNP P30440
A	148	HIS	-	EXPRESSION TAG	UNP P30440
A	149	HIS	-	EXPRESSION TAG	UNP P30440
A	150	HIS	-	EXPRESSION TAG	UNP P30440
A	151	HIS	-	EXPRESSION TAG	UNP P30440
A	152	HIS	-	EXPRESSION TAG	UNP P30440
B	0	MET	-	INITIATING METHIONINE	UNP P30438
B	60	LEU	VAL	CONFLICT	UNP P30438
B	147	HIS	-	EXPRESSION TAG	UNP P30440
B	148	HIS	-	EXPRESSION TAG	UNP P30440
B	149	HIS	-	EXPRESSION TAG	UNP P30440
B	150	HIS	-	EXPRESSION TAG	UNP P30440
B	151	HIS	-	EXPRESSION TAG	UNP P30440
B	152	HIS	-	EXPRESSION TAG	UNP P30440



- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	122	Total	O	0	0
			122	122		
2	B	119	Total	O	0	0
			119	119		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Major allergen I polypeptide, fused chain 1, chain 2



- Molecule 1: Major allergen I polypeptide, fused chain 1, chain 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.59Å 43.00Å 48.91Å 70.80° 80.56° 81.53°	Depositor
Resolution (Å)	36.51 – 1.64 36.48 – 1.64	Depositor EDS
% Data completeness (in resolution range)	92.9 (36.51-1.64) 82.8 (36.48-1.64)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.23 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.191 , 0.220 0.200 , 0.225	Depositor DCC
$R_{free}$ test set	1677 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.7	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 37.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 33376 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2495	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.93	0/1143	1.04	4/1548 (0.3%)
1	B	0.93	0/1139	1.04	5/1543 (0.3%)
All	All	0.93	0/2282	1.04	9/3091 (0.3%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	109	ARG	NE-CZ-NH2	-7.67	116.46	120.30
1	B	130	ASP	CB-CG-OD2	7.44	125.00	118.30
1	A	132	LEU	CB-CG-CD2	6.76	122.49	111.00
1	A	142	ASP	CB-CG-OD2	6.57	124.22	118.30
1	B	9	ASP	CB-CG-OD2	6.24	123.91	118.30
1	B	132	LEU	CB-CG-CD2	6.16	121.47	111.00
1	A	145	GLY	N-CA-C	-6.01	98.07	113.10
1	A	130	ASP	CB-CG-OD2	5.88	123.59	118.30
1	B	46	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1129	0	1145	19	0
1	B	1125	0	1142	19	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	122	0	0	9	1
2	B	119	0	0	6	1
All	All	2495	0	2287	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:ASP:HB3	2:B:261:HOH:O	1.48	1.14
1:A:0:MET:HG3	2:A:249:HOH:O	1.52	1.09
1:A:8:ARG:NH1	1:B:121:GLU:OE2	2.00	0.93
1:B:62:ASP:OD2	2:B:261:HOH:O	2.00	0.79
1:B:124:LEU:HD23	1:B:124:LEU:O	1.86	0.76
1:A:132:LEU:O	1:A:136:THR:HG23	1.87	0.74
1:B:124:LEU:HD21	1:B:128:VAL:HG13	1.74	0.70
1:A:55:GLU:OE2	2:A:246:HOH:O	2.12	0.66
1:B:132:LEU:O	1:B:136:THR:HG23	1.95	0.66
1:A:8:ARG:NE	2:A:226:HOH:O	2.00	0.65
1:B:124:LEU:C	1:B:124:LEU:HD23	2.18	0.64
1:A:89:ASN:ND2	2:A:158:HOH:O	2.32	0.63
1:A:8:ARG:HH12	1:B:121:GLU:CD	2.02	0.60
1:A:142:ASP:HB3	2:A:195:HOH:O	2.01	0.60
1:B:120:VAL:HA	1:B:125:ILE:HG12	1.87	0.56
1:A:93:LEU:O	1:A:97:LEU:HG	2.06	0.56
1:B:62:ASP:CB	2:B:261:HOH:O	2.21	0.55
1:A:142:ASP:CB	2:A:195:HOH:O	2.54	0.55
1:A:89:ASN:HA	1:A:125:ILE:HG23	1.92	0.52
1:B:55:GLU:HG2	2:B:189:HOH:O	2.10	0.52
1:B:24:GLN:HG2	2:B:201:HOH:O	2.09	0.51
1:B:13:PHE:CD2	1:B:14:LEU:HD12	2.46	0.50
1:B:124:LEU:CD2	1:B:128:VAL:HG13	2.41	0.50
1:B:124:LEU:HD21	1:B:128:VAL:CG1	2.43	0.48
1:A:23:GLU:O	1:A:27:GLN:HG2	2.13	0.47
1:A:132:LEU:O	1:A:136:THR:CG2	2.61	0.47
1:A:125:ILE:O	2:A:158:HOH:O	2.20	0.46
1:B:10:VAL:O	1:B:14:LEU:HD13	2.15	0.46
1:A:48:LYS:HE2	1:A:117:ASP:HB2	1.96	0.46
1:A:24:GLN:HA	1:A:27:GLN:HG3	1.99	0.45
1:B:140:SER:OG	1:B:142:ASP:OD1	2.25	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:ARG:HD3	2:A:250:HOH:O	2.17	0.44
1:A:24:GLN:HB3	1:A:24:GLN:HE21	1.57	0.44
1:A:8:ARG:NH2	2:A:226:HOH:O	2.53	0.42
1:B:52:GLU:HG3	2:B:246:HOH:O	2.21	0.40
1:B:124:LEU:C	1:B:124:LEU:CD2	2.89	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:154:HOH:O	2:B:241:HOH:O[1_554]	1.99	0.21

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/153 (94%)	141 (98%)	2 (1%)	1 (1%)	26	7
1	B	143/153 (94%)	141 (99%)	1 (1%)	1 (1%)	26	7
All	All	287/306 (94%)	282 (98%)	3 (1%)	2 (1%)	26	7

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	129	LEU
1	B	77	CYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/137 (95%)	124 (95%)	6 (5%)	33	7
1	B	130/137 (95%)	119 (92%)	11 (8%)	13	1
All	All	260/274 (95%)	243 (94%)	17 (6%)	21	3

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	24	GLN
1	A	27	GLN
1	A	132	LEU
1	A	136	THR
1	A	144	MET
1	B	0	MET
1	B	2	ILE
1	B	24	GLN
1	B	62	ASP
1	B	125	ILE
1	B	127	ARG
1	B	128	VAL
1	B	132	LEU
1	B	136	THR
1	B	142	ASP
1	B	144	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	ASN
1	A	103	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	146/153 (95%)	0.01	4 (2%) 58 56	10, 19, 28, 37	0
1	B	145/153 (94%)	-0.07	2 (1%) 78 78	11, 17, 30, 44	0
All	All	291/306 (95%)	-0.03	6 (2%) 67 67	10, 18, 30, 44	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	GLY	3.9
1	B	144	MET	3.0
1	A	97	LEU	2.8
1	A	144	MET	2.6
1	B	142	ASP	2.3
1	A	103	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.