



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 09:12 PM GMT

PDB ID : 4ZOZ
Title : Crystal structure of the Chaetomium thermophilum Sqt1 bound to the N-terminus of the ribosomal protein L10
Authors : Pausch, P.; Altegoer, F.; Bange, G.
Deposited on : 2015-05-07
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	<div> <div>3%</div> <div>76%</div> <div>6%</div> <div>18%</div> </div>
1	B	480	<div> <div>5%</div> <div>76%</div> <div>6%</div> <div>18%</div> </div>
2	C	28	<div> <div>7%</div> <div>36%</div> <div>• •</div> <div>57%</div> </div>
2	Y	28	<div> <div>7%</div> <div>43%</div> <div>57%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sqt1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	0	0
			2907	1834	494	572	7			
1	B	394	Total	C	N	O	S	0	0	0
			2907	1834	494	572	7			

- Molecule 2 is a protein called 60S ribosomal protein L10-like protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Y	12	Total	C	N	O	S	0	0	0
			106	65	25	14	2			
2	C	12	Total	C	N	O	S	0	0	0
			106	65	25	14	2			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	21	GLY	-	expression tag	UNP G0SEI1
Y	22	SER	-	expression tag	UNP G0SEI1
Y	23	HIS	-	expression tag	UNP G0SEI1
Y	24	HIS	-	expression tag	UNP G0SEI1
Y	25	HIS	-	expression tag	UNP G0SEI1
Y	26	HIS	-	expression tag	UNP G0SEI1
Y	27	HIS	-	expression tag	UNP G0SEI1
Y	28	HIS	-	expression tag	UNP G0SEI1
C	21	GLY	-	expression tag	UNP G0SEI1
C	22	SER	-	expression tag	UNP G0SEI1
C	23	HIS	-	expression tag	UNP G0SEI1
C	24	HIS	-	expression tag	UNP G0SEI1
C	25	HIS	-	expression tag	UNP G0SEI1
C	26	HIS	-	expression tag	UNP G0SEI1
C	27	HIS	-	expression tag	UNP G0SEI1
C	28	HIS	-	expression tag	UNP G0SEI1

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	301	Total 301	O 301	0	0
3	Y	11	Total 11	O 11	0	0
3	B	247	Total 247	O 247	0	0
3	C	6	Total 6	O 6	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	47.50 Å 67.18 Å 72.81 Å 89.92° 108.92° 92.03°	Depositor
Resolution (Å)	48.33 – 1.70 48.33 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.6 (48.33-1.70) 89.8 (48.33-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 1.70 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1685)	Depositor
R, R_{free}	0.191 , 0.216 0.190 , 0.215	Depositor DCC
R_{free} test set	4506 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.9	EDS
Estimated twinning fraction	0.025 for h,-k,-h-l 0.018 for -h,k,-l 0.011 for -h,-k,h+l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 87739 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6591	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/2970	0.56	0/4046
1	B	0.37	0/2970	0.57	0/4046
2	C	0.32	0/108	0.58	0/142
2	Y	0.35	0/108	0.55	0/142
All	All	0.37	0/6156	0.57	0/8376

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	224	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2907	0	2809	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2907	0	2809	23	0
2	C	106	0	109	2	0
2	Y	106	0	109	0	0
3	A	301	0	0	8	1
3	B	247	0	0	10	1
3	C	6	0	0	0	0
3	Y	11	0	0	0	0
All	All	6591	0	5836	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:ARG:NH1	1:A:167:TYR:OH	2.00	0.94
1:B:165:ARG:NH1	1:B:167:TYR:OH	2.06	0.88
1:B:490:GLY:N	3:B:605:HOH:O	2.17	0.78
1:A:225:ASP:N	1:B:330:ARG:HH12	1.86	0.73
1:B:505:GLY:O	3:B:601:HOH:O	2.06	0.72
1:A:222:ASP:OD2	3:A:601:HOH:O	2.10	0.70
1:B:437:ASN:HB2	3:B:821:HOH:O	1.94	0.66
1:A:150:PRO:HG2	1:A:151:ARG:HG2	1.78	0.65
1:A:493:LYS:NZ	1:A:533:ALA:O	2.29	0.65
1:B:222:ASP:OD2	3:B:603:HOH:O	2.15	0.63
1:B:76:LEU:HD11	1:B:151:ARG:HG2	1.80	0.63
2:C:12:CYS:O	2:C:13:LYS:HD2	2.02	0.59
1:B:61:PHE:O	3:B:606:HOH:O	2.18	0.57
1:B:437:ASN:ND2	3:B:604:HOH:O	2.16	0.56
1:A:128:SER:N	3:A:610:HOH:O	2.41	0.54
1:A:499:ARG:HD3	3:A:840:HOH:O	2.09	0.51
1:B:493:LYS:HD3	1:B:495:TRP:CZ2	2.46	0.51
1:A:225:ASP:H	1:B:330:ARG:HH12	1.60	0.50
1:B:223:ALA:O	1:B:226:PRO:HD3	2.12	0.50
1:A:172:PRO:HB2	1:A:176:ALA:H	1.77	0.49
1:A:174:ASN:OD1	1:A:174:ASN:O	2.31	0.48
1:B:326:VAL:HG22	1:B:370:ILE:HD13	1.96	0.47
1:A:440:MET:HB3	1:A:444:LEU:HG	1.97	0.47
1:A:224:SER:HB2	3:A:619:HOH:O	2.15	0.47
1:B:437:ASN:N	1:B:437:ASN:OD1	2.40	0.46
1:B:222:ASP:HB2	3:B:619:HOH:O	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225:ASP:HA	1:B:226:PRO:HD2	1.72	0.45
1:A:284:GLN:NE2	3:A:602:HOH:O	2.20	0.45
1:A:419:GLY:HA2	1:A:422:ALA:O	2.17	0.45
1:A:173:GLN:HB2	3:A:624:HOH:O	2.16	0.45
2:C:12:CYS:O	2:C:13:LYS:HB2	2.18	0.44
1:A:253:LEU:HD21	3:A:639:HOH:O	2.18	0.44
1:B:61:PHE:CZ	1:B:63:GLY:HA3	2.52	0.44
1:B:423:GLU:OE1	3:B:608:HOH:O	2.21	0.44
1:B:295:ARG:NH2	3:B:627:HOH:O	2.51	0.43
1:B:172:PRO:HB2	1:B:176:ALA:H	1.83	0.43
1:B:380:ASN:HB2	1:B:398:THR:OG1	2.17	0.42
1:A:495:TRP:CD1	1:A:531:PHE:HB3	2.56	0.41
1:B:525:ASP:OD1	1:B:525:ASP:N	2.49	0.41
1:A:225:ASP:N	1:A:226:PRO:HD3	2.36	0.41
1:A:415:ARG:HD3	3:A:607:HOH:O	2.19	0.41
1:B:165:ARG:NH2	3:B:620:HOH:O	2.43	0.41
1:A:220:THR:HG23	1:A:231:GLN:HB2	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:826:HOH:O	3:B:801:HOH:O[1_556]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/480 (80%)	373 (97%)	11 (3%)	2 (0%)	34	15
1	B	386/480 (80%)	373 (97%)	11 (3%)	2 (0%)	34	15
2	C	10/28 (36%)	10 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	Y	10/28 (36%)	10 (100%)	0	0	100	100
All	All	792/1016 (78%)	766 (97%)	22 (3%)	4 (0%)	34	15

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	175	GLY
1	B	225	ASP
1	A	173	GLN
1	A	175	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/377 (82%)	308 (100%)	0	100	100
1	B	308/377 (82%)	307 (100%)	1 (0%)	94	92
2	C	10/25 (40%)	9 (90%)	1 (10%)	9	2
2	Y	10/25 (40%)	10 (100%)	0	100	100
All	All	636/804 (79%)	634 (100%)	2 (0%)	94	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	227	SER
2	C	13	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	394/480 (82%)	0.28	16 (4%) 41 44	12, 20, 40, 75	0
1	B	394/480 (82%)	0.46	26 (6%) 22 23	12, 25, 48, 77	0
2	C	12/28 (42%)	0.96	2 (16%) 2 2	29, 36, 50, 51	0
2	Y	12/28 (42%)	0.38	2 (16%) 2 2	22, 26, 45, 47	0
All	All	812/1016 (79%)	0.38	46 (5%) 27 29	12, 22, 45, 77	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	175	GLY	13.3
1	A	175	GLY	11.7
1	A	174	ASN	10.1
1	B	174	ASN	9.2
1	B	225	ASP	9.1
1	B	176	ALA	6.7
1	A	225	ASP	6.5
1	A	173	GLN	6.5
1	B	223	ALA	6.3
1	B	177	LEU	6.2
1	B	227	SER	6.1
1	A	227	SER	5.9
1	A	462	GLY	5.9
1	B	462	GLY	5.8
2	C	12	CYS	5.7
1	A	223	ALA	5.6
1	B	437	ASN	5.2
1	A	224	SER	5.0
2	Y	12	CYS	4.9
1	A	228	ASN	4.7
1	B	228	ASN	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	177	LEU	3.9
1	B	92	ASP	3.9
1	A	226	PRO	3.6
1	A	330	ARG	3.6
1	A	89	ASP	3.6
1	B	91	ASP	3.5
1	B	172	PRO	3.5
1	A	176	ALA	3.3
1	B	295	ARG	3.2
2	C	2	ALA	3.1
1	B	226	PRO	3.0
1	B	224	SER	3.0
1	B	330	ARG	2.8
1	B	130	ASN	2.8
1	B	102	ILE	2.7
1	B	89	ASP	2.6
1	B	173	GLN	2.6
1	A	172	PRO	2.5
1	B	240	THR	2.5
2	Y	13	LYS	2.4
1	B	222	ASP	2.4
1	B	132	ILE	2.2
1	A	295	ARG	2.1
1	B	55	ASN	2.1
1	B	129	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.