



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:31 PM GMT

PDB ID : 3ZPR
Title : Thermostabilised turkey beta1 adrenergic receptor with 4-methyl-2-(piperazin-1-yl) quinoline bound
Authors : Christopher, J.A.; Congreve, M.; Dore, A.S.; Marshall, F.H.; Myszka, D.G.; Brown, J.; Koglin, M.; Tehan, B.; Errey, J.C.; Tate, C.G.; Warne, T.
Deposited on : 2013-03-01
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	Y01	A	402	-	-	-	X
3	Y01	B	401	-	-	-	X
3	Y01	B	402	-	-	-	X
5	2CV	A	502	-	-	-	X
5	2CV	B	502	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-1 ADRENERGIC RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2244	1483	366	375	20			
1	B	298	Total	C	N	O	S	0	0	0
			2370	1559	396	395	20			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	MET	-	EXPRESSION TAG	UNP P07700
A	32	GLY	-	EXPRESSION TAG	UNP P07700
A	369	HIS	-	EXPRESSION TAG	UNP P07700
A	370	HIS	-	EXPRESSION TAG	UNP P07700
A	371	HIS	-	EXPRESSION TAG	UNP P07700
A	372	HIS	-	EXPRESSION TAG	UNP P07700
A	373	HIS	-	EXPRESSION TAG	UNP P07700
A	68	SER	ARG	ENGINEERED MUTATION	UNP P07700
A	90	VAL	MET	ENGINEERED MUTATION	UNP P07700
A	116	LEU	CYS	ENGINEERED MUTATION	UNP P07700
A	227	ALA	TYR	ENGINEERED MUTATION	UNP P07700
A	282	LEU	ALA	ENGINEERED MUTATION	UNP P07700
A	327	ALA	PHE	ENGINEERED MUTATION	UNP P07700
A	338	MET	PHE	ENGINEERED MUTATION	UNP P07700
A	358	ALA	CYS	ENGINEERED MUTATION	UNP P07700
B	31	MET	-	EXPRESSION TAG	UNP P07700
B	32	GLY	-	EXPRESSION TAG	UNP P07700
B	369	HIS	-	EXPRESSION TAG	UNP P07700
B	370	HIS	-	EXPRESSION TAG	UNP P07700
B	371	HIS	-	EXPRESSION TAG	UNP P07700
B	372	HIS	-	EXPRESSION TAG	UNP P07700
B	373	HIS	-	EXPRESSION TAG	UNP P07700
B	68	SER	ARG	ENGINEERED MUTATION	UNP P07700
B	90	VAL	MET	ENGINEERED MUTATION	UNP P07700
B	116	LEU	CYS	ENGINEERED MUTATION	UNP P07700

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	227	ALA	TYR	ENGINEERED MUTATION	UNP P07700
B	282	LEU	ALA	ENGINEERED MUTATION	UNP P07700
B	327	ALA	PHE	ENGINEERED MUTATION	UNP P07700
B	338	MET	PHE	ENGINEERED MUTATION	UNP P07700
B	358	ALA	CYS	ENGINEERED MUTATION	UNP P07700

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Na 2 2	0	0
2	A	2	Total Na 2 2	0	0

- Molecule 3 is CHOLESTEROL HEMISUCCINATE (three-letter code: Y01) (formula: C₃₁H₅₀O₄).



-
- ORTEP diagram of the chemical structure 3WC. The structure consists of a fused bicyclic system, specifically a benzopyridine derivative, linked to a piperidine ring. The atoms are labeled with green text: C2, C4, C5, C6, C8, C10, C11, C12, C13, C15, C20, C22, C24, C26, N10, N11, and N12. The piperidine ring is shown in a chair conformation. The benzopyridine system features a pyridine ring fused to a benzene ring. The piperidine ring is attached to the pyridine ring at the 2-position. The atoms are colored: carbon atoms are grey, nitrogen atoms are blue, and hydrogen atoms are white. The structure is shown with thermal ellipsoids at the 50% probability level. The title '3WC' is centered above the structure.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 17	C 14	N 3	0	0
4	B	1	Total 17	C 14	N 3	0	0

-
- Chemical structure of 2CV (2,6-dichloro-4-vinylphenol) is shown. The structure features a central benzene ring with a vinyl group (-CH=CH₂) at position 4 and two chlorine atoms (-Cl) at positions 2 and 6. The atoms are labeled as follows: C1, C2, C3, C4, C5, C6 for the ring carbons; C7, C8 for the vinyl group; O1, O2 for the hydroxyl groups; and Cl1, Cl2 for the chlorine atoms.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 14 12 1 1	0	0
5	A	1	Total C 7 7	0	0
5	A	1	Total C N O 21 16 1 4	0	0
5	B	1	Total C N O 26 18 1 7	0	0
5	B	1	Total C N O 14 12 1 1	0	0
5	B	1	Total C N O 12 10 1 1	0	0
5	B	1	Total C N O 17 14 1 2	0	0
5	B	1	Total C N O 14 12 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	18	Total O 18 18	0	0
6	B	17	Total O 17 17	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BETA-1 ADRENERGIC RECEPTOR



• Molecule 1: BETA-1 ADRENERGIC RECEPTOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	89.97Å 60.81Å 101.18Å 90.00° 109.20° 90.00°	Depositor
Resolution (Å)	95.55 – 2.70 49.45 – 2.70	Depositor EDS
% Data completeness (in resolution range)	95.7 (95.55-2.70) 95.8 (49.45-2.70)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.224 , 0.266 0.220 , 0.270	Depositor DCC
R_{free} test set	1416 reflections (5.42%)	DCC
Wilson B-factor (Å ²)	58.1	Xtriage
Anisotropy	0.258	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 27537 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4952	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 2CV, Y01, 3WC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	4/2298 (0.2%)	0.66	0/3133
1	B	0.61	1/2425 (0.0%)	0.66	0/3301
All	All	0.62	5/4723 (0.1%)	0.66	0/6434

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	40	TRP	CD2-CE2	6.45	1.49	1.41
1	A	40	TRP	CD2-CE2	6.31	1.49	1.41
1	A	181	TRP	CD2-CE2	5.29	1.47	1.41
1	A	107	TRP	CD2-CE2	5.17	1.47	1.41
1	A	303	TRP	CD2-CE2	5.09	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2244	0	2319	25	0
1	B	2370	0	2468	19	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	70	0	98	6	0
3	B	70	0	98	6	0
4	A	17	0	17	2	0
4	B	17	0	17	1	0
5	A	42	0	59	2	0
5	B	83	0	117	13	0
6	A	18	0	0	0	0
6	B	17	0	0	0	0
All	All	4952	0	5193	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:501:2CV:H601	5:B:501:2CV:C27	1.84	1.08
1:A:159:LYS:HE2	3:A:402:Y01:HAL2	1.36	1.08
3:B:402:Y01:HAC1	3:B:402:Y01:HAE2	1.18	1.07
1:A:205:ARG:NH1	5:B:505:2CV:H361	1.72	1.04
1:A:175:LEU:HD13	1:B:175:LEU:HD22	1.50	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/315 (89%)	272 (98%)	7 (2%)	0	100	100
1	B	296/315 (94%)	288 (97%)	8 (3%)	0	100	100
All	All	575/630 (91%)	560 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/273 (89%)	243 (100%)	1 (0%)	93	98
1	B	258/273 (94%)	256 (99%)	2 (1%)	86	96
All	All	502/546 (92%)	499 (99%)	3 (1%)	90	97

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	329	ASN
1	B	241	ILE
1	B	329	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	HIS
1	A	329	ASN
1	B	286	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	Y01	A	401	-	35,38,38	1.19	3 (8%)	54,57,57	2.34	12 (22%)
3	Y01	A	402	-	35,38,38	1.20	2 (5%)	54,57,57	1.36	7 (12%)
4	3WC	A	500	-	19,19,19	1.85	5 (26%)	26,26,26	1.56	5 (19%)
5	2CV	A	501	-	13,13,25	0.76	0	14,14,30	0.61	0
5	2CV	A	502	-	6,6,25	0.50	0	5,5,30	0.26	0
5	2CV	A	503	-	19,20,25	0.64	0	19,23,30	0.92	1 (5%)
3	Y01	B	401	-	35,38,38	1.09	2 (5%)	54,57,57	1.90	11 (20%)
3	Y01	B	402	-	35,38,38	1.10	2 (5%)	54,57,57	1.38	7 (12%)
4	3WC	B	500	-	19,19,19	1.76	4 (21%)	26,26,26	1.82	7 (26%)
5	2CV	B	501	-	25,25,25	0.52	0	29,30,30	1.09	3 (10%)
5	2CV	B	502	-	13,13,25	0.65	0	14,14,30	0.61	0
5	2CV	B	503	-	11,11,25	0.63	0	11,11,30	0.40	0
5	2CV	B	505	-	16,16,25	0.91	1 (6%)	16,18,30	0.60	0
5	2CV	B	506	-	13,13,25	0.71	0	14,14,30	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	Y01	A	401	-	-	0/17/77/77	0/4/4/4
3	Y01	A	402	-	-	0/17/77/77	0/4/4/4
4	3WC	A	500	-	-	0/4/12/12	0/3/3/3
5	2CV	A	501	-	-	0/13/13/34	0/0/0/0
5	2CV	A	502	-	-	0/4/4/34	0/0/0/0
5	2CV	A	503	-	-	0/24/24/34	0/0/0/0
3	Y01	B	401	-	-	0/17/77/77	0/4/4/4
3	Y01	B	402	-	-	0/17/77/77	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	3WC	B	500	-	-	0/4/12/12	0/3/3/3
5	2CV	B	501	-	-	0/34/34/34	0/0/0/0
5	2CV	B	502	-	-	0/13/13/34	0/0/0/0
5	2CV	B	503	-	-	0/9/9/34	0/0/0/0
5	2CV	B	505	-	-	0/17/17/34	0/0/0/0
5	2CV	B	506	-	-	0/13/13/34	0/0/0/0

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	500	3WC	C6-C5	-5.45	1.40	1.51
3	B	402	Y01	CAK-CAI	-4.90	1.39	1.50
3	A	402	Y01	CAK-CAI	-4.47	1.40	1.50
3	B	401	Y01	CAK-CAI	-4.35	1.40	1.50
4	B	500	3WC	C6-C5	-4.32	1.42	1.51

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	Y01	CAR-CBC-CAV	-7.21	99.12	111.06
3	A	401	Y01	CBC-CAV-CAZ	-4.84	103.59	111.52
3	B	401	Y01	CAR-CBC-CAV	-4.20	104.11	111.06
3	B	402	Y01	CAR-CBC-CAV	-3.94	104.54	111.06
4	B	500	3WC	C2-C22-N11	-3.22	119.76	122.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	Y01	2	0
3	A	402	Y01	4	0
4	A	500	3WC	2	0
5	A	503	2CV	2	0
3	B	401	Y01	2	0
3	B	402	Y01	4	0
4	B	500	3WC	1	0
5	B	501	2CV	5	0
5	B	502	2CV	2	0
5	B	503	2CV	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	505	2CV	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/315 (89%)	0.37	21 (7%) 17 15	33, 53, 96, 144	0
1	B	298/315 (94%)	0.36	22 (7%) 17 15	33, 52, 84, 105	0
All	All	581/630 (92%)	0.36	43 (7%) 17 15	33, 53, 90, 144	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	283	MET	10.1
1	A	282	LEU	9.8
1	A	286	HIS	7.5
1	A	281	MET	6.9
1	B	351	LYS	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	Y01	B	402	35/35	0.91	0.37	4.45	65,71,114,121	0
3	Y01	B	401	35/35	0.95	0.28	4.24	45,52,83,89	0
3	Y01	A	402	35/35	0.88	0.29	3.59	67,85,107,108	0
5	2CV	B	502	14/26	0.87	0.35	2.40	52,79,95,101	0
5	2CV	B	503	12/26	0.83	0.29	1.84	59,72,90,95	0
5	2CV	B	505	17/26	0.76	0.28	1.35	67,79,102,103	0
5	2CV	B	501	26/26	0.81	0.24	1.34	47,69,87,92	0
5	2CV	A	502	7/26	0.92	0.45	1.31	57,64,76,76	0
5	2CV	A	503	21/26	0.90	0.25	0.73	35,60,69,72	0
3	Y01	A	401	35/35	0.96	0.21	0.39	43,50,63,70	0
5	2CV	B	506	14/26	0.82	0.22	0.34	51,72,84,86	0
4	3WC	B	500	17/17	0.96	0.17	-0.35	35,38,40,41	0
2	NA	B	2	1/1	0.39	0.14	-0.43	57,57,57,57	0
5	2CV	A	501	14/26	0.71	0.13	-0.63	90,102,117,120	0
2	NA	A	2	1/1	0.82	0.09	-1.10	57,57,57,57	0
4	3WC	A	500	17/17	0.98	0.16	-1.45	43,47,50,52	0
2	NA	B	1	1/1	0.96	0.10	-2.30	52,52,52,52	0
2	NA	A	1	1/1	0.98	0.06	-5.46	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.