



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:31 PM GMT

PDB ID : 3ZQQ  
Title : Crystal structure of the full-length small terminase from a SPP1-like bacteriophage  
Authors : Buttner, C.R.; Chechik, M.; Ortiz-Lombardia, M.; Smits, C.; Chechik, V.; Jeschke, G.; Dykeman, E.; Benini, S.; Alonso, J.C.; Antson, A.A.  
Deposited on : 2011-06-10  
Resolution : 4.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1010 (4.42-3.56)
Clashscore	102246	1052 (4.40-3.60)
Ramachandran outliers	100387	1005 (4.40-3.60)
Sidechain outliers	100360	1013 (4.42-3.58)
RSRZ outliers	91569	1013 (4.42-3.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	<div> <div>6%</div> <div>41% 32% 7% 20%</div> </div>
1	B	164	<div> <div>%</div> <div>18% 26% .. 53%</div> </div>
1	C	164	<div> <div>13%</div> <div>41% 37% .. 18%</div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2646 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TERMINASE SMALL SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	132	Total	C	N	O	S	0	0	0
			1021	640	184	195	2			
1	B	77	Total	C	N	O	S	0	0	0
			591	370	105	115	1			
1	C	134	Total	C	N	O	S	0	0	0
			1034	648	187	197	2			

There are 57 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	EXPRESSION TAG	UNP Q1EJR8
A	-17	GLY	-	EXPRESSION TAG	UNP Q1EJR8
A	-16	SER	-	EXPRESSION TAG	UNP Q1EJR8
A	-15	SER	-	EXPRESSION TAG	UNP Q1EJR8
A	-14	HIS	-	EXPRESSION TAG	UNP Q1EJR8
A	-13	HIS	-	EXPRESSION TAG	UNP Q1EJR8
A	-12	HIS	-	EXPRESSION TAG	UNP Q1EJR8
A	-11	HIS	-	EXPRESSION TAG	UNP Q1EJR8
A	-10	HIS	-	EXPRESSION TAG	UNP Q1EJR8
A	-9	HIS	-	EXPRESSION TAG	UNP Q1EJR8
A	-8	SER	-	EXPRESSION TAG	UNP Q1EJR8
A	-7	SER	-	EXPRESSION TAG	UNP Q1EJR8
A	-6	GLY	-	EXPRESSION TAG	UNP Q1EJR8
A	-5	LEU	-	EXPRESSION TAG	UNP Q1EJR8
A	-4	VAL	-	EXPRESSION TAG	UNP Q1EJR8
A	-3	PRO	-	EXPRESSION TAG	UNP Q1EJR8
A	-2	ARG	-	EXPRESSION TAG	UNP Q1EJR8
A	-1	GLY	-	EXPRESSION TAG	UNP Q1EJR8
A	0	SER	-	EXPRESSION TAG	UNP Q1EJR8
B	-18	MET	-	EXPRESSION TAG	UNP Q1EJR8
B	-17	GLY	-	EXPRESSION TAG	UNP Q1EJR8
B	-16	SER	-	EXPRESSION TAG	UNP Q1EJR8
B	-15	SER	-	EXPRESSION TAG	UNP Q1EJR8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	EXPRESSION TAG	UNP Q1EJR8
B	-13	HIS	-	EXPRESSION TAG	UNP Q1EJR8
B	-12	HIS	-	EXPRESSION TAG	UNP Q1EJR8
B	-11	HIS	-	EXPRESSION TAG	UNP Q1EJR8
B	-10	HIS	-	EXPRESSION TAG	UNP Q1EJR8
B	-9	HIS	-	EXPRESSION TAG	UNP Q1EJR8
B	-8	SER	-	EXPRESSION TAG	UNP Q1EJR8
B	-7	SER	-	EXPRESSION TAG	UNP Q1EJR8
B	-6	GLY	-	EXPRESSION TAG	UNP Q1EJR8
B	-5	LEU	-	EXPRESSION TAG	UNP Q1EJR8
B	-4	VAL	-	EXPRESSION TAG	UNP Q1EJR8
B	-3	PRO	-	EXPRESSION TAG	UNP Q1EJR8
B	-2	ARG	-	EXPRESSION TAG	UNP Q1EJR8
B	-1	GLY	-	EXPRESSION TAG	UNP Q1EJR8
B	0	SER	-	EXPRESSION TAG	UNP Q1EJR8
C	-18	MET	-	EXPRESSION TAG	UNP Q1EJR8
C	-17	GLY	-	EXPRESSION TAG	UNP Q1EJR8
C	-16	SER	-	EXPRESSION TAG	UNP Q1EJR8
C	-15	SER	-	EXPRESSION TAG	UNP Q1EJR8
C	-14	HIS	-	EXPRESSION TAG	UNP Q1EJR8
C	-13	HIS	-	EXPRESSION TAG	UNP Q1EJR8
C	-12	HIS	-	EXPRESSION TAG	UNP Q1EJR8
C	-11	HIS	-	EXPRESSION TAG	UNP Q1EJR8
C	-10	HIS	-	EXPRESSION TAG	UNP Q1EJR8
C	-9	HIS	-	EXPRESSION TAG	UNP Q1EJR8
C	-8	SER	-	EXPRESSION TAG	UNP Q1EJR8
C	-7	SER	-	EXPRESSION TAG	UNP Q1EJR8
C	-6	GLY	-	EXPRESSION TAG	UNP Q1EJR8
C	-5	LEU	-	EXPRESSION TAG	UNP Q1EJR8
C	-4	VAL	-	EXPRESSION TAG	UNP Q1EJR8
C	-3	PRO	-	EXPRESSION TAG	UNP Q1EJR8
C	-2	ARG	-	EXPRESSION TAG	UNP Q1EJR8
C	-1	GLY	-	EXPRESSION TAG	UNP Q1EJR8
C	0	SER	-	EXPRESSION TAG	UNP Q1EJR8





## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.18 Å 98.18 Å 323.82 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	107.00 – 4.00 107.94 – 4.01	Depositor EDS
% Data completeness (in resolution range)	61.5 (107.00-4.00) 61.9 (107.94-4.01)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.83 (at 4.01 Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, $R_{free}$	0.200 , 0.282 0.218 , 0.278	Depositor DCC
$R_{free}$ test set	481 reflections (14.55%)	DCC
Wilson B-factor (Å <sup>2</sup> )	110.1	Xtriage
Anisotropy	1.281	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 224.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 3307 reflections	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	2646	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	219.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1029	0.43	0/1378
1	B	0.22	0/594	0.46	0/796
1	C	0.22	0/1042	0.44	0/1394
All	All	0.24	0/2665	0.44	0/3568

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

There are no bond length outliers.

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	132	THR	CB

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1021	0	1065	68	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	591	0	621	57	0
1	C	1034	0	1081	58	0
All	All	2646	0	2767	141	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:LEU:HD21	1:B:68:LEU:HD11	1.39	1.05
1:B:111:ILE:HD11	1:C:80:ILE:HG21	1.58	0.84
1:A:129:LYS:HB3	1:B:127:LYS:HG2	1.63	0.80
1:B:77:LEU:HA	1:B:80:ILE:HD12	1.64	0.79
1:B:137:ILE:HD11	1:C:135:VAL:HG22	1.63	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/164 (78%)	111 (87%)	14 (11%)	3 (2%)	8	51
1	B	75/164 (46%)	62 (83%)	10 (13%)	3 (4%)	4	37
1	C	130/164 (79%)	107 (82%)	18 (14%)	5 (4%)	4	39
All	All	333/492 (68%)	280 (84%)	42 (13%)	11 (3%)	5	43

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	133	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	67	ILE
1	C	6	LEU
1	C	68	LEU
1	A	140	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/134 (80%)	95 (89%)	12 (11%)	7	37
1	B	64/134 (48%)	59 (92%)	5 (8%)	16	54
1	C	108/134 (81%)	104 (96%)	4 (4%)	41	75
All	All	279/402 (69%)	258 (92%)	21 (8%)	17	56

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	ILE
1	A	128	GLN
1	C	46	LEU
1	A	75	GLU
1	C	54	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	121	HIS
1	C	88	GLN
1	C	45	ASN
1	A	128	GLN
1	B	128	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

### 5.7 Other polymers ⓘ

There are no such residues in this entry.

### 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	132/164 (80%)	0.54	10 (7%) 17 12	146, 209, 252, 319	0
1	B	77/164 (46%)	0.53	1 (1%) 79 70	145, 197, 240, 310	0
1	C	134/164 (81%)	0.86	21 (15%) 3 3	140, 226, 339, 383	0
All	All	343/492 (69%)	0.66	32 (9%) 11 8	140, 208, 315, 383	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	33	TYR	5.9
1	C	31	ALA	5.0
1	C	17	TYR	4.5
1	C	51	ILE	4.3
1	C	35	LYS	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.