



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:11 AM GMT

PDB ID : 2ZU6
Title : crystal structure of the eIF4A-PDCD4 complex
Authors : Cho, Y.; Chang, J.H.; Sohn, S.Y.
Deposited on : 2008-10-13
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<div> <div>3%</div> <div> <div>45%</div> <div>35%</div> <div>9%</div> <div>12%</div> </div> </div>
1	C	388	<div> <div>9%</div> <div> <div>40%</div> <div>40%</div> <div>11%</div> <div>10%</div> </div> </div>
1	D	388	<div> <div>3%</div> <div> <div>47%</div> <div>33%</div> <div>9%</div> <div>11%</div> </div> </div>
1	F	388	<div> <div>18%</div> <div> <div>31%</div> <div>31%</div> <div>7%</div> <div>30%</div> </div> </div>
2	B	307	<div> <div>0%</div> <div> <div>56%</div> <div>32%</div> <div>7%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	307	<div> <div></div> <div>%</div> <div>49%</div> <div>36%</div> <div>7%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	3	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic initiation factor 4A-I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	343	Total	C	N	O	S	0	0	0
			2743	1734	477	514	18			
1	C	351	Total	C	N	O	S	0	0	0
			2819	1778	492	531	18			
1	D	345	Total	C	N	O	S	0	0	0
			2756	1742	480	516	18			
1	F	271	Total	C	N	O	S	0	0	0
			2196	1389	381	411	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	MET	-	EXPRESSION TAG	UNP P60842
C	19	MET	-	EXPRESSION TAG	UNP P60842
D	19	MET	-	EXPRESSION TAG	UNP P60842
F	19	MET	-	EXPRESSION TAG	UNP P60842

- Molecule 2 is a protein called Programmed cell death protein 4.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	284	Total	C	N	O	S	Se	0	0	0
			2218	1403	366	434	6	9			
2	E	282	Total	C	N	O	S	Se	0	0	0
			2201	1393	363	430	6	9			

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	11	Total	O	0	0
			11	11		
5	B	6	Total	O	0	0
			6	6		
5	C	3	Total	O	0	0
			3	3		
5	D	11	Total	O	0	0
			11	11		
5	E	2	Total	O	0	0
			2	2		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Eukaryotic initiation factor 4A-I






• Molecule 1: Eukaryotic initiation factor 4A-I



• Molecule 1: Eukaryotic initiation factor 4A-I



• Molecule 2: Programmed cell death protein 4

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	156.89Å 165.38Å 100.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 49.86 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.00-2.80) 94.1 (49.86-2.81)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.81Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.248 , 0.294 0.239 , 0.246	Depositor DCC
R_{free} test set	3053 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	62.1	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 63.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 64546 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14986	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.77	2/2781 (0.1%)	0.84	4/3750 (0.1%)
1	C	0.41	0/2855	0.54	0/3846
1	D	0.72	0/2791	0.81	1/3761 (0.0%)
1	F	0.44	0/2218	0.54	0/2978
2	B	0.65	0/2244	0.73	1/3015 (0.0%)
2	E	0.67	1/2227 (0.0%)	0.75	1/2993 (0.0%)
All	All	0.63	3/15116 (0.0%)	0.71	7/20343 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1
2	E	0	2
All	All	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	192	PHE	N-CA	7.02	1.60	1.46
2	E	227	CYS	CB-SG	-5.21	1.73	1.81
1	A	193	LYS	CE-NZ	-5.07	1.36	1.49

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	PHE	CB-CG-CD1	-8.22	115.05	120.80
2	E	334	LEU	CA-CB-CG	6.31	129.82	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	192	PHE	N-CA-C	5.92	126.97	111.00
1	A	22	VAL	CB-CA-C	-5.53	100.90	111.40
1	A	155	ILE	CG1-CB-CG2	-5.16	100.05	111.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	E	286	VAL	Peptide
2	E	316	GLY	Peptide
1	F	152	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2743	0	2792	160	0
1	C	2819	0	2865	202	0
1	D	2756	0	2811	136	0
1	F	2196	0	2220	122	0
2	B	2218	0	2231	80	0
2	E	2201	0	2214	117	0
3	A	8	0	12	1	0
3	D	4	0	6	0	0
4	A	4	0	3	1	0
4	D	4	0	3	0	0
5	A	11	0	0	0	0
5	B	6	0	0	0	0
5	C	3	0	0	1	0
5	D	11	0	0	1	0
5	E	2	0	0	1	0
All	All	14986	0	15157	782	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 782 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:279:ILE:HD13	1:C:284:LYS:HB3	1.25	1.15
1:F:115:GLN:O	1:F:119:VAL:HG23	1.48	1.14
2:E:442:GLN:NE2	2:E:442:GLN:H	1.45	1.14
2:B:339:LEU:HD11	2:B:370:MSE:HE3	1.29	1.13
1:C:73:ILE:HD12	1:C:224:THR:HG21	1.22	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/388 (86%)	297 (89%)	31 (9%)	7 (2%)	9	29
1	C	339/388 (87%)	265 (78%)	60 (18%)	14 (4%)	3	11
1	D	333/388 (86%)	288 (86%)	32 (10%)	13 (4%)	4	12
1	F	247/388 (64%)	198 (80%)	37 (15%)	12 (5%)	3	8
2	B	280/307 (91%)	250 (89%)	26 (9%)	4 (1%)	14	42
2	E	278/307 (91%)	226 (81%)	44 (16%)	8 (3%)	6	19
All	All	1812/2166 (84%)	1524 (84%)	230 (13%)	58 (3%)	5	17

5 of 58 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	253	VAL
1	A	291	LYS
1	A	296	ASP
2	B	270	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/342 (89%)	251 (83%)	52 (17%)	2	7
1	C	314/342 (92%)	256 (82%)	58 (18%)	2	6
1	D	305/342 (89%)	249 (82%)	56 (18%)	2	6
1	F	245/342 (72%)	177 (72%)	68 (28%)	0	1
2	B	251/259 (97%)	223 (89%)	28 (11%)	7	22
2	E	249/259 (96%)	215 (86%)	34 (14%)	4	13
All	All	1667/1886 (88%)	1371 (82%)	296 (18%)	2	6

5 of 296 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	382	ARG
1	D	243	LEU
1	F	312	ASP
1	C	396	GLU
1	D	123	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 64 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	206	ASN
1	C	374	ASN
1	F	201	GLN
1	C	248	GLN
1	C	280	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACY	A	1	-	1,3,3	1.84	0	0,3,3	0.00	-
3	EDO	A	2	-	3,3,3	0.54	0	2,2,2	0.33	0
3	EDO	A	3	-	3,3,3	0.60	0	2,2,2	0.27	0
3	EDO	D	1	-	3,3,3	0.54	0	2,2,2	0.57	0
4	ACY	D	2	-	1,3,3	1.29	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACY	A	1	-	-	0/0/0/0	0/0/0/0
3	EDO	A	2	-	-	0/1/1/1	0/0/0/0
3	EDO	A	3	-	-	0/1/1/1	0/0/0/0
3	EDO	D	1	-	-	0/1/1/1	0/0/0/0
4	ACY	D	2	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1	ACY	1	0
3	A	3	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	343/388 (88%)	0.23	12 (3%) 48 35	30, 64, 115, 161	0
1	C	351/388 (90%)	0.64	35 (9%) 9 4	56, 125, 160, 163	0
1	D	345/388 (88%)	0.16	11 (3%) 51 39	31, 68, 116, 160	0
1	F	271/388 (69%)	1.26	68 (25%) 1 0	69, 122, 155, 161	0
2	B	275/307 (89%)	-0.01	2 (0%) 89 84	36, 65, 90, 108	0
2	E	273/307 (88%)	-0.03	3 (1%) 82 74	37, 70, 92, 108	0
All	All	1858/2166 (85%)	0.37	131 (7%) 19 10	30, 84, 149, 163	0

The worst 5 of 131 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	48	TYR	7.9
1	F	175	TYR	6.9
1	C	389	THR	5.5
1	F	212	LEU	5.4
1	F	331	LEU	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	A	3	4/4	0.89	0.32	3.91	57,62,64,66	0
3	EDO	D	1	4/4	0.91	0.24	1.48	67,68,69,69	0
4	ACY	A	1	4/4	0.83	0.22	0.41	78,78,79,79	0
4	ACY	D	2	4/4	0.83	0.37	-	75,75,76,76	0
3	EDO	A	2	4/4	0.81	0.22	-	81,83,84,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.