Full wwPDB X-ray Structure Validation Report (i)

Jun 9, 2016 – 06:15 AM EDT

PDB ID	:	4ZUH
Title	:	Complex structure of PEDV 3CLpro mutant (C144A) with a peptide substrate.
Authors	:	Ye, G.; Fu, Z.F.; Peng, G.Q.
Deposited on	:	2015-05-16
Resolution	:	2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027674
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027674

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	5%, 86%, 11%
1	B	305	3%, 89%, 8%
2	C	11	36%, 55%, 18%, 27%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEDV 3C-Like protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	2269	1432	392	430	15	0	0	0
1	B	294	2243	1418	388	422	15	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	144	ALA	CYS	engineered mutation	UNP K4L9I6
A	300	HIS	-	expression tag	UNP K4L9I6
A	301	HIS	-	expression tag	UNP K4L9I6
A	302	HIS	-	expression tag	UNP K4L9I6
A	303	HIS	-	expression tag	UNP K4L9I6
A	304	HIS	-	expression tag	UNP K4L9I6
A	305	HIS	-	expression tag	UNP K4L9I6
B	144	ALA	CYS	engineered mutation	UNP K4L9I6
B	300	HIS	-	expression tag	UNP K4L9I6
B	301	HIS	-	expression tag	UNP K4L9I6
B	302	HIS	-	expression tag	UNP K4L9I6
B	303	HIS	-	expression tag	UNP K4L9I6
B	304	HIS	-	expression tag	UNP K4L9I6
B	305	HIS	-	expression tag	UNP K4L9I6

- Molecule 2 is a protein called peptide substrate SAVLQSGF.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	8	56	36	9	11	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	130	130	130	0	0
			Total	O		
3	B	137	137	137	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PEDV 3C-Like protease

Chain A:



- Molecule 1: PEDV 3C-Like protease

Chain B:



- Molecule 2: peptide substrate SAVLQSGF

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	56.68Å 91.88Å 58.01Å 90.00° 100.24° 90.00°	Depositor
Resolution (Å)	31.78 – 2.39 28.70 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (31.78-2.39) 99.4 (28.70-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.79 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.155 , 0.219 0.159 , 0.164	Depositor DCC
R_{free} test set	1140 reflections (4.94%)	DCC
Wilson B-factor (Å^2)	27.5	Xtriage
Anisotropy	0.599	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 49.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4835	wwPDB-VP
Average B, all atoms (Å^2)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.34% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.40	0/2315	0.57	0/3144
1	B	0.41	0/2288	0.54	0/3105
2	C	0.36	0/56	0.56	0/74
All	All	0.41	0/4659	0.56	0/6323

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2269	0	2223	20	0
1	B	2243	0	2198	14	0
2	C	56	0	57	2	0
3	A	130	0	0	4	0
3	B	137	0	0	3	0
All	All	4835	0	4478	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:ARG:HD3	1:A:136:ARG:HE	1.55	0.71
1:A:46:SER:O	3:A:401:HOH:O	2.14	0.65
1:B:217:TRP:CE2	1:B:275:LYS:HD3	2.31	0.65
1:A:132:ASN:ND2	1:A:193:GLU:OE2	2.33	0.61
1:A:148:GLY:HA3	1:A:160:TYR:HB3	1.82	0.61
1:A:31:TRP:CE2	1:A:94:ASN:HB2	2.43	0.54
1:A:296:MET:SD	3:A:407:HOH:O	2.59	0.53
1:B:217:TRP:CD2	1:B:275:LYS:HD3	2.45	0.52
1:A:201:GLU:HG3	3:A:411:HOH:O	2.11	0.50
1:A:40:ARG:HA	1:A:86:LEU:HG	1.95	0.49
1:B:148:GLY:HA3	1:B:160:TYR:HB3	1.96	0.47
1:A:162:HIS:NE2	2:C:5:GLN:OE1	2.31	0.47
1:A:180:MET:CE	1:A:186:ASP:HB3	2.44	0.47
1:A:130:ARG:HD3	1:A:136:ARG:NE	2.28	0.46
1:A:27:LEU:HD21	1:A:42:VAL:HB	1.98	0.46
1:B:230:ASN:HB3	1:B:240:THR:HG22	1.97	0.46
1:A:31:TRP:CD2	1:A:94:ASN:HB2	2.51	0.46
1:B:259:GLN:OE1	3:B:401:HOH:O	2.20	0.46
1:A:114:LEU:HD11	1:A:121:ALA:HB1	1.98	0.45
1:B:292:VAL:O	1:B:296:MET:HG2	2.17	0.44
1:A:1:ALA:HA	3:A:485:HOH:O	2.18	0.43
1:A:112:ASN:O	1:A:148:GLY:HA2	2.18	0.43
1:A:223:ARG:NH2	1:A:257:ASP:OD2	2.41	0.43
1:B:21:CYS:HB2	1:B:66:SER:HB2	2.01	0.43
1:A:12:VAL:CG2	1:A:98:PRO:HB3	2.48	0.43
1:B:40:ARG:HA	1:B:86:LEU:HG	2.00	0.42
1:B:80:THR:HG23	3:B:509:HOH:O	2.20	0.41
1:A:24:ASN:O	2:C:8:PHE:HA	2.20	0.41
1:B:170:CYS:HB3	3:B:525:HOH:O	2.20	0.41
1:B:112:ASN:O	1:B:148:GLY:HA2	2.21	0.41
1:B:31:TRP:CE2	1:B:94:ASN:HB2	2.56	0.41
1:B:217:TRP:CZ2	1:B:275:LYS:HB3	2.56	0.41
1:A:21:CYS:HB2	1:A:66:SER:HB3	2.03	0.40
1:B:180:MET:CE	1:B:186:ASP:HB3	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/305 (97%)	284 (96%)	10 (3%)	2 (1%)	26	38
1	B	290/305 (95%)	287 (99%)	3 (1%)	0	100	100
2	C	6/11 (54%)	5 (83%)	1 (17%)	0	100	100
All	All	592/621 (95%)	576 (97%)	14 (2%)	2 (0%)	46	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	46	SER
1	A	48	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/251 (97%)	236 (97%)	8 (3%)	45	66
1	B	240/251 (96%)	235 (98%)	5 (2%)	61	80
2	C	6/9 (67%)	6 (100%)	0	100	100
All	All	490/511 (96%)	477 (97%)	13 (3%)	52	73

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	60	LEU
1	A	136	ARG
1	A	190	LEU
1	A	196	SER
1	A	267	SER
1	A	271	ASN
1	A	284	THR
1	B	27	LEU
1	B	60	LEU
1	B	80	THR
1	B	99	LYS
1	B	269	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	298/305 (97%)	-0.02	14 (4%) 35 36	13, 26, 52, 86	1 (0%)
1	B	294/305 (96%)	-0.23	8 (2%) 58 57	14, 24, 41, 55	0
2	C	8/11 (72%)	1.80	4 (50%) 0 0	33, 42, 53, 54	0
All	All	600/621 (96%)	-0.10	26 (4%) 39 40	13, 25, 48, 86	1 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	195	ALA	6.9
1	A	48	THR	6.5
2	C	8	PHE	4.6
1	A	47	THR	4.0
1	A	46	SER	3.8
1	A	153	ASN	3.5
1	A	242	VAL	3.3
1	B	167	GLY	3.3
1	B	188	PRO	3.1
1	B	192	VAL	3.1
2	C	1	SER	3.0
1	B	153	ASN	3.0
1	B	195	ALA	2.9
1	A	194	GLY	2.8
1	A	84	ALA	2.6
2	C	2	ALA	2.5
1	A	192	VAL	2.4
1	A	61	ARG	2.3
1	A	45	SER	2.1
1	A	152	ASN	2.1
2	C	7	GLY	2.1
1	A	63	HIS	2.1
1	B	154	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	189	THR	2.0
1	A	83	GLY	2.0
1	B	168	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.