Full wwPDB X-ray Structure Validation Report (i)

Feb 1, 2016 – 12:08 AM GMT

PDB ID	:	1ZVT
Title	:	Structure of the E. coli ParC C-terminal domain
Authors	:	Corbett, K.D.; Schoeffler, A.J.; Thomsen, N.D.; Berger, J.M.
Deposited on	:	2005-06-02
Resolution	:	1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	2% 87% 9% •
1	B	256	5% 89% 7% •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Topoisomerase IV subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1849	1159	331	349	10	0	0	0
1	B	246	1856	1164	332	350	10	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	209	209	209	0	0
2	B	171	171	171	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Topoisomerase IV subunit A

Chain A:



- Molecule 1: Topoisomerase IV subunit A

Chain B:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	40.91Å 50.49Å 72.76Å 86.11° 86.90° 70.57°	Depositor
Resolution (Å)	20.00 – 1.70 29.61 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.2 (20.00-1.70) 94.7 (29.61-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	3.52 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.183 , 0.215 0.185 , 0.217	Depositor DCC
R_{free} test set	2937 reflections (5.34%)	DCC
Wilson B-factor (Å^2)	20.8	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 45.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 57955 reflections	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4085	wwPDB-VP
Average B, all atoms (Å^2)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.85% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.53	0/1879	0.67	1/2540 (0.0%)
1	B	0.49	0/1888	0.66	0/2555
All	All	0.51	0/3767	0.66	1/5095 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	692	LEU	CA-CB-CG	5.32	127.53	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1849	0	1900	15	0
1	B	1856	0	1908	12	0
2	A	209	0	0	2	0
2	B	171	0	0	4	0
All	All	4085	0	3808	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:740:ASP:HB2	2:B:244:HOH:O	1.71	0.90
1:A:681:ARG:NH1	1:A:683:GLU:OE1	2.09	0.86
1:B:642:MET:HE3	1:B:654:MET:HG2	1.71	0.72
1:A:541:GLN:HG3	2:A:340:HOH:O	1.90	0.71
1:A:710:ARG:HB3	1:A:711:PRO:HD2	1.86	0.57
1:B:642:MET:CE	1:B:654:MET:HG2	2.36	0.55
1:B:742:PRO:C	2:B:173:HOH:O	2.46	0.54
1:A:711:PRO:O	1:A:715:GLN:HG3	2.07	0.53
1:B:670:LYS:NZ	1:B:673:ASN:HD22	2.07	0.53
1:A:705:ARG:HD3	2:A:233:HOH:O	2.11	0.51
1:A:693:PRO:HG2	1:A:741:SER:HB2	1.94	0.49
1:A:556:ASP:OD1	1:A:558:ILE:HG13	2.13	0.49
1:A:700:ILE:HG21	1:A:731:LEU:HD11	1.95	0.48
1:B:670:LYS:HZ2	1:B:673:ASN:HD22	1.62	0.47
1:A:736:ARG:NH2	1:B:688:GLN:OE1	2.49	0.45
1:A:611:LEU:HG	1:A:621:LEU:HD22	1.99	0.44
1:A:710:ARG:HB3	1:A:711:PRO:CD	2.46	0.44
1:B:705:ARG:NH1	2:B:378:HOH:O	2.51	0.44
1:A:509:TRP:CE2	1:A:526:LYS:HE2	2.53	0.43
1:B:701:HIS:HB2	1:B:736:ARG:HB2	2.00	0.42
1:A:741:SER:HA	1:A:742:PRO:HD3	1.88	0.41
1:B:687:ALA:C	1:B:688:GLN:HG2	2.40	0.41
1:A:642:MET:HE3	1:A:654:MET:HG2	2.02	0.41
1:A:556:ASP:CG	1:A:558:ILE:HG13	2.41	0.41
1:B:511:ARG:HD2	1:B:563:ALA:HA	2.03	0.40
1:B:532:LYS:NZ	2:B:357:HOH:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	241/256 (94%)	240 (100%)	1 (0%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	244/256 (95%)	239 (98%)	5 (2%)	0	100	100
All	All	485/512 (95%)	479 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/210 (96%)	199 (99%)	2 (1%)	82	72
1	B	202/210 (96%)	197 (98%)	5 (2%)	55	34
All	All	403/420 (96%)	396 (98%)	7 (2%)	68	51

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	692	LEU
1	A	728	MET
1	B	518	ILE
1	B	590	ASP
1	B	593	LYS
1	B	692	LEU
1	B	729	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	524	ASN
1	A	648	GLN
1	A	673	ASN
1	A	688	GLN
1	B	516	HIS
1	B	609	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	648	GLN
1	B	673	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	245/256 (95%)	-0.01	5 (2%)	68	73	16, 22, 33, 40	0
1	B	246/256 (96%)	0.22	14 (5%)	27	29	17, 25, 41, 48	0
All	All	491/512 (95%)	0.11	19 (3%)	43	47	16, 24, 37, 48	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	528	GLY	4.9
1	B	564	ARG	4.0
1	B	566	GLN	3.8
1	A	497	SER	3.6
1	B	517	ASP	3.3
1	B	527	ALA	3.3
1	B	733	ARG	3.2
1	A	564	ARG	3.0
1	B	514	LYS	2.8
1	A	742	PRO	2.6
1	B	704	LYS	2.5
1	B	498	GLU	2.5
1	A	704	LYS	2.5
1	B	742	PRO	2.3
1	B	578	PRO	2.2
1	B	730	GLY	2.2
1	B	524	ASN	2.1
1	B	545	PHE	2.1
1	A	527	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.