



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 27, 2016 – 05:22 AM EDT

PDB ID : 4ZVQ
Title : Caspase-7 Variant 2 (V2) with reprogrammed substrate specificity due to Y230V/W232M/Q276C substitutions bound to VEID inhibitor.
Authors : Hill, M.E.; MacPherson, D.J.; Hardy, J.A.
Deposited on : 2015-05-18
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027790
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	<div> <div>65%</div> <div>5%</div> <div>30%</div> </div>
1	C	198	<div> <div>64%</div> <div>6%</div> <div>29%</div> </div>
2	B	113	<div> <div>4%</div> <div>79%</div> <div>18%</div> </div>
2	D	113	<div> <div>4%</div> <div>73%</div> <div>9%</div> <div>18%</div> </div>
3	E	5	<div> <div>80%</div> <div>20%</div> </div>
3	F	5	<div> <div>80%</div> <div>20%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	0	0
			1092	687	187	207	11			
1	C	140	Total	C	N	O	S	0	0	0
			1099	691	188	209	11			

- Molecule 2 is a protein called Caspase-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	93	Total	C	N	O	S	0	0	0
			755	484	127	138	6			
2	D	93	Total	C	N	O	S	0	0	0
			755	484	127	138	6			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	230	VAL	TYR	engineered mutation	UNP P55210
B	232	MET	TRP	engineered mutation	UNP P55210
B	276	CYS	GLN	engineered mutation	UNP P55210
B	304	LEU	-	expression tag	UNP P55210
B	305	GLU	-	expression tag	UNP P55210
B	306	HIS	-	expression tag	UNP P55210
B	307	HIS	-	expression tag	UNP P55210
B	308	HIS	-	expression tag	UNP P55210
B	309	HIS	-	expression tag	UNP P55210
B	310	HIS	-	expression tag	UNP P55210
B	311	HIS	-	expression tag	UNP P55210
D	530	VAL	TYR	engineered mutation	UNP P55210
D	532	MET	TRP	engineered mutation	UNP P55210
D	576	CYS	GLN	engineered mutation	UNP P55210
D	604	LEU	-	expression tag	UNP P55210
D	605	GLU	-	expression tag	UNP P55210

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	606	HIS	-	expression tag	UNP P55210
D	607	HIS	-	expression tag	UNP P55210
D	608	HIS	-	expression tag	UNP P55210
D	609	HIS	-	expression tag	UNP P55210
D	610	HIS	-	expression tag	UNP P55210
D	611	HIS	-	expression tag	UNP P55210

- Molecule 3 is a protein called Peptide ACE-VAL-GLU-ILE-ASA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	5	Total	C	N	O	0	0	0
			35	22	4	9			
3	F	5	Total	C	N	O	0	0	0
			35	22	4	9			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	29	Total	O	0	0
			29	29		
4	B	16	Total	O	0	0
			16	16		
4	C	34	Total	O	0	0
			34	34		
4	D	19	Total	O	0	0
			19	19		
4	F	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Caspase-7




• Molecule 1: Caspase-7




• Molecule 2: Caspase-7



• Molecule 2: Caspase-7




• Molecule 3: Peptide ACE-VAL-GLU-ILE-ASA





- Molecule 3: Peptide ACE-VAL-GLU-ILE-ASA

Chain F:  80% 20%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	86.89Å 86.89Å 187.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	69.85 – 2.50 69.85 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (69.85-2.50) 99.9 (69.85-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.175 , 0.208 0.176 , 0.208	Depositor DCC
R_{free} test set	1413 reflections (4.85%)	DCC
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.475	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3870	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ASA, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/1109	0.45	0/1486
1	C	0.25	0/1116	0.48	1/1496 (0.1%)
2	B	0.23	0/775	0.44	0/1047
2	D	0.23	0/775	0.46	0/1047
3	E	0.20	0/24	0.69	0/32
3	F	0.21	0/24	0.61	0/32
All	All	0.24	0/3823	0.46	1/5140 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	422	LEU	CA-CB-CG	6.34	129.87	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1092	0	1078	6	0
1	C	1099	0	1085	8	0
2	B	755	0	736	3	0
2	D	755	0	736	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	35	0	32	1	0
3	F	35	0	32	1	0
4	A	29	0	0	0	0
4	B	16	0	0	0	0
4	C	34	0	0	0	0
4	D	19	0	0	0	0
4	F	1	0	0	0	0
All	All	3870	0	3699	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:463:THR:HG21	2:D:521:PHE:HE2	1.56	0.70
2:D:533:ARG:HA	2:D:539:SER:HA	1.78	0.65
2:B:233:ARG:HA	2:B:239:SER:HA	1.87	0.57
1:C:397:LEU:HD13	1:C:440:ILE:HG21	1.86	0.56
1:A:196:GLN:HE21	2:D:513:ILE:HD12	1.71	0.56
1:A:163:THR:HG21	2:B:221:PHE:HE1	1.70	0.55
2:B:275:SER:HB2	3:E:1:VAL:HG11	1.92	0.51
1:A:97:LEU:HD13	1:A:140:ILE:HG21	1.93	0.51
1:C:371:ILE:HG21	1:C:422:LEU:HD21	1.94	0.49
2:D:575:SER:HB2	3:F:1:VAL:HG11	1.95	0.48
1:C:474:LEU:HA	1:C:477:LYS:HD2	1.95	0.48
1:C:476:GLU:HG2	2:D:513:ILE:HD13	1.96	0.46
1:C:495:ILE:O	1:C:496:GLN:HB3	2.16	0.44
1:C:493:ASP:OD1	2:D:585:LYS:NZ	2.31	0.43
2:D:546:CYS:O	2:D:550:GLU:HG3	2.18	0.42
1:A:174:LEU:HA	1:A:177:LYS:HD2	2.00	0.42
1:C:384:MET:HB3	1:C:444:HIS:CD2	2.55	0.41
1:A:119:MET:O	1:A:123:LEU:HD23	2.21	0.41
1:A:70:CYS:HA	1:A:138:ALA:O	2.21	0.40
2:D:566:ASN:OD1	2:D:589:PRO:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/198 (69%)	135 (98%)	2 (2%)	0	100	100
1	C	138/198 (70%)	135 (98%)	3 (2%)	0	100	100
2	B	91/113 (80%)	90 (99%)	1 (1%)	0	100	100
2	D	91/113 (80%)	90 (99%)	1 (1%)	0	100	100
3	E	3/5 (60%)	3 (100%)	0	0	100	100
3	F	3/5 (60%)	3 (100%)	0	0	100	100
All	All	463/632 (73%)	456 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/172 (69%)	119 (100%)	0	100	100
1	C	120/172 (70%)	120 (100%)	0	100	100
2	B	85/103 (82%)	85 (100%)	0	100	100
2	D	85/103 (82%)	85 (100%)	0	100	100
3	E	3/3 (100%)	3 (100%)	0	100	100
3	F	3/3 (100%)	3 (100%)	0	100	100
All	All	415/556 (75%)	415 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	196	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ASA	E	4	1,3	2,7,7	0.82	0	2,8,8	1.44	1 (50%)
3	ASA	F	4	1,3	2,7,7	0.83	0	2,8,8	1.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASA	E	4	1,3	-	0/2/6/6	0/0/0/0
3	ASA	F	4	1,3	-	0/2/6/6	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	4	ASA	O-C-CA	-2.01	120.34	125.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	139/198 (70%)	-0.14	0	100 100	30, 40, 55, 69	0
1	C	140/198 (70%)	-0.11	0	100 100	27, 35, 51, 75	0
2	B	93/113 (82%)	0.05	4 (4%)	39 44	27, 38, 78, 91	0
2	D	93/113 (82%)	0.16	4 (4%)	39 44	26, 37, 77, 88	0
3	E	3/5 (60%)	0.04	0	100 100	47, 47, 54, 57	0
3	F	3/5 (60%)	0.61	0	100 100	44, 44, 46, 50	0
All	All	471/632 (74%)	-0.03	8 (1%)	73 76	26, 38, 68, 91	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	580	PRO	4.6
2	D	578	ASP	2.9
2	B	211	TYR	2.8
2	B	280	PRO	2.5
2	D	576	CYS	2.4
2	B	281	HIS	2.2
2	B	278	ASP	2.1
2	D	582	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ASA	E	4	8/8	0.99	0.12	-	38,42,43,44	0
3	ASA	F	4	8/8	0.98	0.11	-	33,37,42,44	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.