



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:11 AM GMT

PDB ID : 2ZWA
Title : Crystal structure of tRNA wybutosine synthesizing enzyme TYW4
Authors : Suzuki, Y.; Noma, A.; Suzuki, T.; Ishitani, R.; Nureki, O.
Deposited on : 2008-12-01
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CIT	B	802	-	-	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 12533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Leucine carboxyl methyltransferase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	672	Total 5413	C 3470	N 899	O 1018	S 26	0	8	0
1	B	683	Total 5475	C 3507	N 915	O 1027	S 26	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	417	LEU	MET	SEE REMARK 999	UNP Q08282
B	417	LEU	MET	SEE REMARK 999	UNP Q08282


- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$).



Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	B	1	26	14	6	5	1	0	0

- Molecule 3 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	13	6	7	0	0
3	B	1	13	6	7	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	811	Total O 811 811	0	0
5	B	732	Total O 732 732	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leucine carboxyl methyltransferase 2



- Molecule 1: Leucine carboxyl methyltransferase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.58Å 89.44Å 236.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.62 – 1.70 38.62 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.7 (38.62-1.70) 98.4 (38.62-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 1.71Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.170 , 0.211 0.168 , 0.209	Depositor DCC
R_{free} test set	17318 reflections (9.91%)	DCC
Wilson B-factor (Å ²)	21.3	Xtrriage
Anisotropy	0.272	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.5	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	1 of 174757 reflections (0.001%)	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	12533	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, EDO, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/5552	0.56	0/7506
1	B	0.37	0/5599	0.56	0/7569
All	All	0.37	0/11151	0.56	0/15075

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5413	0	5432	72	0
1	B	5475	0	5500	68	0
2	A	26	0	19	0	0
2	B	26	0	19	0	0
3	A	13	0	5	0	0
3	B	13	0	5	0	0
4	A	12	0	18	0	0
4	B	12	0	18	0	0
5	A	811	0	0	12	0
5	B	732	0	0	9	0
All	All	12533	0	11016	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:606:THR:HG22	1:A:608:ARG:H	1.21	1.02
1:B:102:GLN:HE22	1:B:355:ALA:H	0.90	0.88
1:A:592:HIS:HD2	1:A:594:LEU:H	1.16	0.88
1:A:102:GLN:HE22	1:A:355:ALA:H	1.23	0.85
1:B:396:ALA:HB3	1:B:475:MET:HE1	1.60	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	674/695 (97%)	655 (97%)	19 (3%)	0	100	100
1	B	680/695 (98%)	663 (98%)	16 (2%)	1 (0%)	56	35
All	All	1354/1390 (97%)	1318 (97%)	35 (3%)	1 (0%)	56	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	407	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	616/630 (98%)	599 (97%)	17 (3%)	51	29
1	B	622/630 (99%)	607 (98%)	15 (2%)	57	36
All	All	1238/1260 (98%)	1206 (97%)	32 (3%)	54	32

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	606	THR
1	B	11	ASN
1	B	517	LEU
1	A	682	SER
1	B	95	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	592	HIS
1	B	102	GLN
1	B	416	GLN
1	A	685	ASN
1	B	137	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	696	-	3,3,3	0.52	0	2,2,2	0.37	0
4	EDO	A	697	-	3,3,3	0.65	0	2,2,2	0.27	0
4	EDO	A	698	-	3,3,3	0.45	0	2,2,2	0.56	0
2	SAH	A	801	-	20,28,28	1.00	2 (10%)	19,40,40	3.31	7 (36%)
3	CIT	A	802	-	3,12,12	1.23	0	3,17,17	1.61	1 (33%)
4	EDO	B	696	-	3,3,3	0.45	0	2,2,2	0.43	0
4	EDO	B	697	-	3,3,3	0.57	0	2,2,2	0.31	0
4	EDO	B	698	-	3,3,3	0.50	0	2,2,2	0.56	0
2	SAH	B	801	-	20,28,28	1.05	2 (10%)	19,40,40	3.04	3 (15%)
3	CIT	B	802	-	3,12,12	1.10	0	3,17,17	1.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	696	-	-	0/1/1/1	0/0/0/0
4	EDO	A	697	-	-	0/1/1/1	0/0/0/0
4	EDO	A	698	-	-	0/1/1/1	0/0/0/0
2	SAH	A	801	-	-	0/7/31/31	0/3/3/3
3	CIT	A	802	-	-	0/6/16/16	0/0/0/0
4	EDO	B	696	-	-	0/1/1/1	0/0/0/0
4	EDO	B	697	-	-	0/1/1/1	0/0/0/0
4	EDO	B	698	-	-	0/1/1/1	0/0/0/0
2	SAH	B	801	-	-	0/7/31/31	0/3/3/3
3	CIT	B	802	-	-	0/6/16/16	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	SAH	C2-N1	2.23	1.38	1.33
2	A	801	SAH	C2-N1	2.32	1.38	1.33
2	A	801	SAH	C2-N3	3.38	1.38	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	SAH	C2-N3	3.61	1.38	1.32

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	SAH	N3-C2-N1	-12.31	119.47	128.89
2	B	801	SAH	N3-C2-N1	-11.87	119.81	128.89
2	A	801	SAH	C2'-C1'-N9	-5.01	106.63	114.29
2	B	801	SAH	C5'-SD-CG	-3.47	91.99	102.41
2	B	801	SAH	C1'-N9-C4	-2.51	123.15	126.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	672/695 (96%)	0.07	30 (4%) 37 41	12, 23, 51, 81	0
1	B	683/695 (98%)	0.09	28 (4%) 41 44	12, 26, 54, 75	0
All	All	1355/1390 (97%)	0.08	58 (4%) 39 43	12, 24, 54, 81	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	421	TYR	10.9
1	B	421	TYR	7.4
1	B	29	GLY	6.4
1	B	176	TYR	5.7
1	B	358	GLN	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CIT	B	802	13/13	0.95	0.14	7.18	21,28,34,42	0
3	CIT	A	802	13/13	0.92	0.13	1.06	24,32,51,54	0
4	EDO	A	698	4/4	0.93	0.13	1.00	24,27,31,31	0
4	EDO	B	696	4/4	0.99	0.10	0.86	17,19,21,21	0
4	EDO	B	698	4/4	0.93	0.12	0.63	34,34,41,41	0
2	SAH	A	801	26/26	0.97	0.10	0.43	13,17,19,21	0
4	EDO	B	697	4/4	0.97	0.08	-0.19	22,22,24,26	0
4	EDO	A	697	4/4	0.94	0.11	-0.28	23,24,24,27	0
2	SAH	B	801	26/26	0.92	0.09	-0.49	23,29,35,43	0
4	EDO	A	696	4/4	0.98	0.08	-0.57	19,19,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.