



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 21, 2016 – 06:22 PM EDT

PDB ID : 4ZW2  
Title : Crystal structure of the Mouse voltage gated calcium channel beta subunit isoform 1a in complex with Alpha Interaction Domain peptide.  
Authors : Norris, N.C.; Oakley, A.J.  
Deposited on : 2015-05-19  
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	<div> <div>9%</div> <div>88%</div> <div>7% . .</div> </div>
2	B	20	<div> <div>90%</div> <div>5% 5%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Voltage-dependent L-type calcium channel subunit beta-1, Voltage-dependent L-type calcium channel subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	9	0
			2536	1620	441	466	9			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q8R3Z5
A	2	PRO	-	expression tag	UNP Q8R3Z5
A	3	GLY	-	expression tag	UNP Q8R3Z5
A	4	SER	-	expression tag	UNP Q8R3Z5
A	123	HIS	-	linker	UNP Q8R3Z5
A	124	GLY	-	linker	UNP Q8R3Z5
A	125	LYS	-	linker	UNP Q8R3Z5
A	126	ARG	-	linker	UNP Q8R3Z5
A	127	ILE	-	linker	UNP Q8R3Z5
A	128	PRO	-	linker	UNP Q8R3Z5
A	129	PHE	-	linker	UNP Q8R3Z5
A	130	PHE	-	linker	UNP Q8R3Z5
A	131	LYS	-	linker	UNP Q8R3Z5


- Molecule 2 is a protein called Voltage-dependent L-type calcium channel subunit alpha-1S.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	20	Total	C	N	O	S	0	3	1
			172	107	29	34	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	ACE	-	acetylation	UNP Q02789
B	19	NH2	-	amidation	UNP Q02789

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		
3	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	265	Total	O	0	0
			265	265		
4	B	20	Total	O	0	0
			20	20		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Voltage-dependent L-type calcium channel subunit beta-1, Voltage-dependent L-type calcium channel subunit beta-1



- Molecule 2: Voltage-dependent L-type calcium channel subunit alpha-1S



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.24Å 69.37Å 131.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.81 – 1.86 22.81 – 1.86	Depositor EDS
% Data completeness (in resolution range)	99.7 (22.81-1.86) 99.8 (22.81-1.86)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.74 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.169 , 0.207 0.180 , 0.218	Depositor DCC
$R_{free}$ test set	1801 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.8	Xtriage
Anisotropy	0.195	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3010	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.96	2/2617 (0.1%)	0.97	3/3552 (0.1%)
2	B	1.31	2/180 (1.1%)	1.40	4/240 (1.7%)
All	All	0.99	4/2797 (0.1%)	1.01	7/3792 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	8[A]	ARG	CZ-NH1	6.50	1.41	1.33
2	B	8[B]	ARG	CZ-NH1	6.50	1.41	1.33
1	A	299	GLU	CD-OE1	5.56	1.31	1.25
1	A	333	SER	CA-CB	5.14	1.60	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	8[A]	ARG	NE-CZ-NH2	-9.17	115.72	120.30
2	B	8[B]	ARG	NE-CZ-NH2	-9.17	115.72	120.30
1	A	103	ASP	CB-CG-OD1	6.62	124.26	118.30
2	B	8[A]	ARG	NE-CZ-NH1	6.42	123.51	120.30
2	B	8[B]	ARG	NE-CZ-NH1	6.42	123.51	120.30
1	A	117	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	A	140	ASP	CB-CG-OD2	-5.05	113.76	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2536	0	2552	13	0
2	B	172	0	168	5	0
3	A	17	0	23	0	0
4	A	265	0	0	5	0
4	B	20	0	0	4	0
All	All	3010	0	2743	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:8[B]:ARG:CG	4:B:301:HOH:O	2.15	0.94
2:B:8[B]:ARG:HG3	4:B:301:HOH:O	1.78	0.82
1:A:211:THR:HG23	4:A:549:HOH:O	1.89	0.72
1:A:110[A]:GLU:HG3	4:A:624:HOH:O	1.91	0.70
1:A:36:LYS:NZ	1:A:75:GLU:OE1	2.24	0.63
2:B:8[B]:ARG:NE	4:B:301:HOH:O	1.91	0.59
1:A:233:GLN:HG2	4:A:503:HOH:O	2.07	0.54
1:A:61:VAL:O	1:A:61:VAL:HG12	2.09	0.52
2:B:8[B]:ARG:HG2	4:B:301:HOH:O	1.98	0.51
1:A:268:LEU:HD23	1:A:287:ILE:HD13	1.96	0.48
1:A:184[A]:ARG:NH2	1:A:240:ASP:OD2	2.47	0.47
1:A:269[B]:GLN:HG3	1:A:287:ILE:HD11	1.98	0.46
1:A:110[A]:GLU:CG	4:A:624:HOH:O	2.56	0.45
2:B:8[A]:ARG:HG3	2:B:8[A]:ARG:HH11	1.80	0.45
1:A:68:LYS:HA	1:A:68:LYS:HD2	1.80	0.43
1:A:132:LYS:NZ	4:A:502:HOH:O	2.26	0.41
1:A:228:LEU:HD23	1:A:228:LEU:HA	1.89	0.41
1:A:55[B]:GLU:HA	1:A:55[B]:GLU:OE1	2.21	0.41

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/333 (98%)	320 (98%)	6 (2%)	0	100	100
2	B	21/20 (105%)	21 (100%)	0	0	100	100
All	All	347/353 (98%)	341 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/296 (93%)	267 (97%)	9 (3%)	45	25
2	B	19/16 (119%)	18 (95%)	1 (5%)	28	10
All	All	295/312 (95%)	285 (97%)	10 (3%)	46	24

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	27	LEU
1	A	29	GLN
1	A	68	LYS
1	A	206	ILE
1	A	209	SER
1	A	212	ARG
1	A	269[A]	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	269[B]	GLN
2	B	1	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	284	ASN
1	A	295	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PGE	A	401	-	9,9,9	0.69	0	8,8,8	0.49	0
3	PGE	A	402	-	6,6,9	0.51	0	5,5,8	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	401	-	-	0/7/7/7	0/0/0/0
3	PGE	A	402	-	-	0/4/4/7	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/333 (95%)	0.50	29 (9%) <span>11</span> <span>11</span>	26, 38, 66, 111	6 (1%)
2	B	18/20 (90%)	0.21	0 <span>100</span> <span>100</span>	28, 32, 50, 54	0
All	All	337/353 (95%)	0.49	29 (8%) <span>13</span> <span>12</span>	26, 37, 66, 111	6 (1%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	15	ARG	6.9
1	A	186	THR	6.1
1	A	17	ALA	5.2
1	A	199	ASN	5.1
1	A	19	ARG	5.1
1	A	18	LEU	5.1
1	A	133	THR	5.0
1	A	16	GLU	4.8
1	A	127	ILE	4.6
1	A	128	PRO	4.3
1	A	132	LYS	4.2
1	A	22	ALA	4.1
1	A	195	SER	3.8
1	A	26	ALA	3.7
1	A	20	LYS	3.5
1	A	21	GLU	3.4
1	A	96	ILE	3.2
1	A	120	SER	3.2
1	A	198	ASN	3.2
1	A	197	LEU	3.0
1	A	123	HIS	2.9
1	A	33	ALA	2.7
1	A	271	LEU	2.7
1	A	135	HIS	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	200	PRO	2.6
1	A	134	GLU	2.4
1	A	202	LYS	2.3
1	A	71	LEU	2.2
1	A	210	ASN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	PGE	A	401	10/10	0.85	0.17	-	57,63,74,74	0
3	PGE	A	402	7/10	0.78	0.24	-	62,74,76,77	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.